

FluxCalR: a R package for calculating CO₂ and CH₄ fluxes from static chambers

Junbin Zhao^{1, 2}

1 Department of Biogeochemistry and Soil Quality, Division of Environment and Natural Resources, Norwegian Institute of Bioeconomy Research, Ås, Norway **2** Department of Biological Sciences, Florida International University, Miami, FL, USA

DOI: [10.21105/joss.01751](https://doi.org/10.21105/joss.01751)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Arfon Smith](#) ↗

Reviewers:

- [@jhollist](#)
- [@ha0ye](#)

Submitted: 13 August 2019

Published: 21 November 2019

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC-BY](#)).

Summary

As the main drivers of climate change, greenhouse gas (e.g., CO₂ and CH₄) emissions have been monitored intensively across the globe. The static chamber is one of the most commonly used approaches for measuring greenhouse gas fluxes from ecosystems (e.g., stem/soil respiration, CH₄ emission, etc.) because of its easy implementation, high accuracy and low cost (Pumpanen et al., 2004). To perform the measurements, a gas analyzer is usually used to measure the changes of greenhouse gas concentrations within a closed chamber that covers an area of interest (e.g., soil surface) over a certain period of time (usually several minutes). The flux rates (F) are then calculated from the recorded gas concentrations assuming that the changing rate is linear:

$$F = vol / (R \cdot Ta \cdot area) \cdot dG / dt$$

where vol is the volume of the chamber (l), R is the universal gas constant (l atm K⁻¹ mol⁻¹), Ta is the ambient temperature (K), $area$ is the area of the chamber base (m²), and dG/dt is the rate of the measured gas concentration change over time t (ppm s⁻¹) (i.e., the slope of the linear regression).

Even though the method itself appears to be straightforward, to calculate the fluxes from the raw gas concentration data can be tedious, because one usually has to manually extract the data for each measurement from an irregular mix of valid data and unwanted records. After selecting the valid data, it is important to choose a window that delivers the best regression for each measurement to ensure the quality and representativeness of the calculated flux. The R^2 of the regression is usually taken to determine the quality of a flux measurement and a regression with low R^2 is usually discarded (Zhao, Peichl, & Nilsson, 2016, 2017). Therefore, a tool that calculates the gas concentration changes into fluxes in an efficient and objective fashion is needed.

There are some programs that calculate the fluxes measured using static chamber method, such as the Trace Gas Analyzer Platform (LI-COR Inc., Lincoln, NE, USA), the R package `flux` (Jurasinski, Koebisch, Guenther, & Beetz, 2014), and Flux Puppy (Carbone et al., 2019). However, they are either compatible to only one designated system, unable to calculate fluxes for multiple gases, or hard to handle continuously measured gas concentrations. Therefore there is still a need for a free, flexible, and easy-to-use software that is designed to calculate ecosystem gas fluxes from continuously measured gas concentrations.

This R package, FluxCalR is designed to calculate CO₂ and CH₄ gas fluxes measured with static chamber method. It provides an easy solution to calculate all the fluxes in one single data file exported by most gas analyzers that includes continuously measured gas concentrations

for multiple measurements. Overall, users can complete the flux calculation following three simple steps that implement the build-in functions: 1) load the raw dataset and convert it to the required format; 2) define the time cues that correspond to the start and/or end of the measurements, and 3) calculate the fluxes based on the time cues.

FluxCa1R has several features to facilitate the process of flux calculation. Particularly, it has multiple options to identify valid data for the measurements which greatly improves the flexibility and efficiency of flux calculation. It is also flexible in the inputs (e.g., ambient temperature, gas type) that can meet the needs of measurements with different protocols. It ensures the best quality and representativeness of each flux by choosing the regression with the greatest determination coefficient (R^2) within the defined data range, making the calculated results more reproducible. In addition, the FluxCa1R also provides graphs to visualize the linear regressions on top of the original data to ensure no mismatch occurs.

FluxCa1R is meant to be used by, for example, ecosystem researchers who carry out greenhouse gas flux measurements using the static chamber method. It simplifies and shortens the process of tedious flux calculations and helps one focus more on data analysis with the ready calculated fluxes. The functions in the package can be integrated into users' data processing/analyzing workflow. At the same time, it also saves the calculated fluxes to a local file that can be loaded by R again later or further processed in other programs. The original functions in the package have been used for flux calculations in the previous publications (Zhao et al., 2016, 2017) and in other ongoing research projects performed in the Everglades.

Acknowledgements

This work is partially supported by the U.S. Department of Energy's (DOE) National Institute for Climate Change Research (07-SC-NICCR-1059) and National Science Foundation Division of Atmospheric & Geospace Sciences Atmospheric Chemistry Program (1561161, 1561139, 1233006 and 1807533). I also acknowledge the suggestions from Sparkle L. Malone and the assistance of Matthew J. Simon and Michal Gazovic for testing the code on different datasets.

References

- Carbone, M. S., Seyednasrollah, B., Rademacher, T. T., Basler, D., Moine, J. M. L., Beals, S., Beasley, J., et al. (2019). Flux puppy – an open-source software application and portable system design for low-cost manual measurements of CO₂ and H₂O fluxes. *Agricultural and Forest Meteorology*, 274, 1–6. doi:<https://doi.org/10.1016/j.agrformet.2019.04.012>
- Jurasinski, G., Koepsch, F., Guenther, A., & Beetz, S. (2014). Flux: Flux rate calculation from dynamic closed chamber measurements. Retrieved from <https://CRAN.R-project.org/package=flux>
- Pumpanen, J., Kolari, P., Ilvesniemi, H., Minkkinen, K., Vesala, T., Niinistö, S., Lohila, A., et al. (2004). Comparison of different chamber techniques for measuring soil CO₂ efflux. *Agricultural and Forest Meteorology*, 123(3), 159–176. doi:<https://doi.org/10.1016/j.agrformet.2003.12.001>
- Zhao, J., Peichl, M., & Nilsson, M. B. (2016). Enhanced winter soil frost reduces methane emission during the subsequent growing season in a boreal peatland. *Global Change Biology*, 22(2), 750–762. doi:[10.1111/gcb.13119](https://doi.org/10.1111/gcb.13119)
- Zhao, J., Peichl, M., & Nilsson, M. B. (2017). Long-term enhanced winter soil frost alters growing season CO₂ fluxes through its impact on vegetation development in a boreal peatland. *Global Change Biology*, 1–15. doi:[10.1111/gcb.13621](https://doi.org/10.1111/gcb.13621)