

catwoman: A transit modelling Python package for asymmetric light curves

Kathryn Jones¹ and Néstor Espinoza²

¹ University of Bern, Center for Space and Habitability, Gesellschaftsstrasse 6, CH-3012, Bern, Switzerland ² Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

DOI: [10.21105/joss.02382](https://doi.org/10.21105/joss.02382)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Juanjo Bazán](#) ↗

Reviewers:

- [@jessie-dotson](#)
- [@barentsens](#)

Submitted: 12 June 2020

Published: 17 November 2020

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

When exoplanets pass in front of their stars from our point of view on Earth, they imprint a transit signature on the stellar light curve which, to date, has been assumed to be symmetric in time, owing to the planet being modelled as a circular area occulting the stellar surface ([Kreidberg, 2015](#); [Luger et al., 2019](#); see, e.g., [Mandel & Agol, 2002](#)). However this signature might be asymmetric due to several possible effects, one of which is the different temperature/pressure and/or chemical compositions the different terminator regions a transiting planet could have (see, e.g., [Powell et al., 2019](#)). Being able to model these asymmetric signatures directly from transit light curves could give us an unprecedented glimpse into planetary 3-dimensional structure, helping constrain models of atmospheric evolution, structure and composition.

`catwoman` is a Python package that models these asymmetric transit light curves, calculating light curves for any radially symmetric stellar limb darkening law and where planets are modelled as two semi-circles, of different radii, using the integration algorithm developed in ([Kreidberg, 2015](#)) and implemented in the `batman` library, from which `catwoman` builds upon. It is fast and efficient and open source with full documentation available to view at <https://catwoman.readthedocs.io>.

The light curves are modelled as follows: The decrease in flux, δ , as a planet transits its star can be approximated by the sum

$$\delta = \sum_{i=1}^N I(x_m) \Delta A(x_m, R_{p,1}, R_{p,2}, \varphi, d), \quad (1)$$

splitting the semi-circles into iso-intensity bands centred on the star and for each intersectional segment (see [Figure 1](#)) you multiply its area, ΔA , by the intensity of the star and then sum these strips to generate the full δ for a specific separation between the centre of the star and planet, d . The code then increments d by a small pre-determined amount (based on the time array given by the user) and recalculates δ .




Figure 1: Diagram of the geometric configuration during transit of two stacked semi-circles (one of radius $R_{p,1}$, and another of radius $R_{p,2}$) that model the different limbs of an exoplanet transiting in front of a star. The area of the star has been divided in different sections of radius x_i (dashed circles) — between each subsequent section, the star is assumed to have a radially symmetric intensity profile (e.g., blue band between x_{i-1} and x_i above). In order to obtain the light curve, the challenge is to calculate the sum of the intersectional areas between a given iso-intensity band and the semi-circles, ΔA (blue band with dashed grey lines). Note the stacked semi-circles are inclined by an angle φ with respect to the planetary orbital motion.

The width of the iso-intensity bands determines the truncation error of the model. The model is first initialised with parameters including a maximum truncation error either set by the user or taken as the pre-set value as 1ppm. As in `batman`, `catwoman` first calculates many models, with varying widths and geometrically searches for a width that produces an error less than 1% away (and always less than) the specified level. The model then uses this width value to calculate the desired light curves. A lower specified error, and therefore thinner iso-intensity bands, produces more accurate light curves, however more steps are needed to calculate δ which takes more time.

`catwoman` also allows for φ , the angle of rotation of the semi-circles, to vary as a free parameter, which is something no other model has tried to implement, accounting for the possibility of spin-orbit misalignments of the planet. The two semi-circle radii, $R_{p,1}$ and $R_{p,2}$, and other orbital variables are also completely free parameters.

`catwoman` was designed to be used by astronomical researchers. For a realistic light curve with 100 in-transit data points, `catwoman` takes around 340 seconds to produce 1 million quadratic-limb-darkened light curves on a single 1.3 GHz Intel Core i5 processor. It is used in Espinoza & Jones (in prep.).

Acknowledgements

We would like to thank the Max Plank Institute of Astronomy, Heidelberg, for providing the funding for this project and hosting Kathryn Jones as a summer student at the Institute.

References

- Kreidberg, L. (2015). batman: BAsic Transit Model cAlculatioN in Python. *Publications of the ASP*, 127(957), 1161. <https://doi.org/10.1086/683602>
- Luger, R., Agol, E., Foreman-Mackey, D., Fleming, D. P., Lustig-Yaeger, J., & Deitrick, R. (2019). starry: Analytic Occultation Light Curves. *Astronomical Journal*, 157(2), 64. <https://doi.org/10.3847/1538-3881/aae8e5>
- Mandel, K., & Agol, E. (2002). Analytic Light Curves for Planetary Transit Searches. *Astrophysical Journal, Letters*, 580(2), L171–L175. <https://doi.org/10.1086/345520>
- Powell, D., Louden, T., Kreidberg, L., Zhang, X., Gao, P., & Parmentier, V. (2019). Transit Signatures of Inhomogeneous Clouds on Hot Jupiters: Insights from Microphysical Cloud Modeling. *Astrophysical Journal*, 887(2), 170. <https://doi.org/10.3847/1538-4357/ab55d9>