

AstroPaint: A Python Package for Painting Halo Catalogs into Celestial Maps

Siavash Yasini^{*1}, Marcelo Alvarez^{2, 3}, Emmanuel Schaan^{2, 3}, Karime Maamari^{1, 5}, Shobeir K. S. Mazinani⁴, Nareg Mirzatury¹, and Elena Pierpaoli¹

1 University of Southern California 2 Lawrence Berkeley National Laboratory 3 University of California, Berkeley 4 Aetna Inc. 5 Argonne National Lab

DOI: [10.21105/joss.02608](https://doi.org/10.21105/joss.02608)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Alice Harpole](#) ↗

Reviewers:

- [@AshKelly](#)
- [@zpace](#)

Submitted: 13 August 2020

Published: 05 November 2020

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Overview

AstroPaint is a python package for generating and visualizing sky maps of a wide range of astrophysical signals originating from dark matter halos or the gas that they host. AstroPaint creates a whole-sky mock map of the target signal/observable, at a desired resolution, by combining an input halo catalog and the radial/angular profile of the astrophysical effect (see the workflow section for details). The package also provides a suite of tools that can facilitate analysis routines such as catalog filtering, map manipulation, and cutout stacking. The simulation suite has an Object-Oriented design and runs in parallel, making it both easy to use and readily scalable for production of high resolution maps with large underlying catalogs. Although the package has been primarily developed to simulate signals pertinent to galaxy clusters, its application extends to halos of arbitrary size or even point sources.




Figure 1: Map of the Birkinshaw-Gull effect painted with AstroPaint using the WebSky catalog

*corresponding author

Statement of Need

Studying the large scale structure of the universe heavily relies on observations of astrophysical signals at various frequencies. Examples of such studies include detection or characterization of objects such as galaxies, clusters, or voids through either gravitational lensing, electromagnetic scattering, absorption or emission events in the optical, radio, or x-ray frequency bands. Such studies typically require simulated high resolution maps of various astrophysical effects to emulate both the signal and noise (foregrounds) components. For example, in a study that aims to evaluate the detection significance of the Birkinshaw-Gull (BG) effect – a probe of the transverse velocities of halos (Birkinshaw & Gull, 1983; Yasini et al., 2019) – using the Simons Observatory (Abitbol & others, 2019) or CMB-S4 (Abazajian & others, 2019), one needs a mock map of the BG effect Figure 1 as well as maps of potential contaminants such as kinetic and thermal Sunyaev-Zeldovich effects (kSZ and tSZ) (Sunyaev & Zeldovich, 1970) for the same set of objects.

While it is possible to create realistic maps of astrophysical effects through hydrodynamical simulations (Dolag et al., 2016), these methods are numerically expensive for large numbers of objects and reproducing them under different cosmologies and initial conditions can be prohibitive. An alternative strategy for creating mock observations of extended objects such as galaxies and galaxy cluster halos is to simulate the positions of these objects (either semi-analytically or through N-body simulations (Sehgal et al., 2010; Stein et al., 2019; Stein et al., 2020)) and then synthetically paint the desired signal at the location of the halos. AstroPaint is developed to help researchers in creating mock maps using the latter strategy.

AstroPaint can also be used to create templates for detecting astrophysical effects in image data. For example, to detect kSZ for an ensemble of galaxies in a CMB map, one needs a template of this effect for the observed patch of the sky. Such a template can be generated by taking the catalog of the target galaxies along with their velocities and painting kSZ profiles around them on a map using AstroPaint.

Package Structure and Workflow

AstroPaint consists of three main objects that interact with each other: `Catalog`, `Canvas`, and `Painter`.

`Catalog` contains the locations, velocities, and masses of the objects. `Canvas` contains the map of the astrophysical signal in HEALPix format (Zonca et al., 2019). `Painter` contains the template for the radial profile of the signal to be painted on the `Canvas` in circular discs centered at the location of the halos in the `Catalog`.

These objects are sequentially passed into each other according to the following workflow:

```
from astropaint import Catalog, Canvas, Painter

catalog = Catalog(data=input_data)
canvas = Canvas(catalog, nside)
painter = Painter(template=radial_profile)

painter.spray(canvas)
```

The output map array can be accessed via `canvas.pixels` or directly visualized using `canvas.show_map()`. Here `input_data` is the dataframe that hold the locations, velocities, and masses of the halos. `nside` is a parameter in `healpy` that determines the total number of pixels (`npix = 12 * nside ** 2`) and consequently the resolution of the map. Finally,

`radial_profile` is a one-dimensional function that determines the shape of the profile. A mind map visualization of the package structure can be found in [here](#).

Acknowledgements

We would like to thank Simone Ferraro, Tim Morton, Vera Gluscevic, George Stein, Mathew Madhavacheril, Zack Li, and Alex van Engelen for their incredibly helpful comments and feedback. SY is grateful to the BCCP group at UC Berkeley for their hospitality during Summer 2019 where this project was inaugurated. EP is supported by NASA 80NSSC18K0403 and the Simons Foundation award number 615662; EP and SY are supported by NSF AST-1910678.

References

- Abazajian, K., & others. (2019). CMB-S4 Decadal Survey APC White Paper. *Bull. Am. Astron. Soc.*, 51(7), 209. <https://doi.org/10.2172/1556957>
- Abitbol, M. H., & others. (2019). The Simons Observatory: Astro2020 Decadal Project Whitepaper. *Bull. Am. Astron. Soc.*, 51, 147. <http://arxiv.org/abs/1907.08284>
- Birkinshaw, M., & Gull, S. F. (1983). *A test for transverse motions of clusters of galaxies*. 302(5906), 315–317. <https://doi.org/10.1038/302315a0>
- Dolag, K., Komatsu, E., & Sunyaev, R. (2016). SZ effects in the Magneticum Pathfinder Simulation: Comparison with the Planck, SPT, and ACT results. *Mon. Not. Roy. Astron. Soc.*, 463(2), 1797–1811. <https://doi.org/10.1093/mnras/stw2035>
- Sehgal, N., Bode, P., Das, S., Hernand ez-Monteagudo, C., Huffenberger, K., Lin, Y.-T., Ostriker, J. P., & Trac, H. (2010). *Simulations of the Microwave Sky*. 709(2), 920–936. <https://doi.org/10.1088/0004-637X/709/2/920>
- Stein, G., Alvarez, M. A., & Bond, J. R. (2019). The mass-Peak Patch algorithm for fast generation of deep all-sky dark matter halo catalogues and its N-Body validation. *Mon. Not. Roy. Astron. Soc.*, 483(2), 2236–2250. <https://doi.org/10.1093/mnras/sty3226>
- Stein, G., Alvarez, M. A., Bond, J. R., van Engelen, A., & Battaglia, N. (2020). The Websky Extragalactic CMB Simulations. *arXiv E-Prints*, arXiv:2001.08787. <https://doi.org/10.1088/1475-7516/2020/10/012>
- Sunyaev, R. A., & Zeldovich, Y. (1970). The Interaction of matter and radiation in the hot model of the universe. *Astrophys. Space Sci.*, 7, 20–30.
- Yasini, S., Mirzatumy, N., & Pierpaoli, E. (2019). Pairwise Transverse Velocity Measurement with the Rees–Sciama Effect. *Astrophys. J. Lett.*, 873(2), L23. <https://doi.org/10.3847/2041-8213/ab0bfe>
- Zonca, A., Singer, L., Lenz, D., Reinecke, M., Rosset, C., Hivon, E., & Gorski, K. (2019). healpy: equal area pixelization and spherical harmonics transforms for data on the sphere in Python. *Journal of Open Source Software*, 4(35), 1298. <https://doi.org/10.21105/joss.01298>