

# PyAstroPol: A Python package for the instrumental polarization analysis of the astronomical optics.

Hemanth Pruthvi.<sup>1</sup>

<sup>1</sup> Leibniz-institut für Sonnenphysik, Freiburg, Germany.

DOI: [10.21105/joss.02693](https://doi.org/10.21105/joss.02693)

## Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

---

Editor: [Amy Roberts](#) ↗

## Reviewers:

- [@aquilesC](#)
- [@caldarolamartin](#)
- [@mw craig](#)

Submitted: 25 August 2020

Published: 24 November 2020

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## Statement of Need

The instrumental polarization analysis is one of the key aspects of the optical system analysis of the astronomical telescopes and instruments. The majority of the optical surfaces in the astronomical instruments induce a change in the state of polarization of the incoming light, which could lead to inaccurate measurements of the state of polarization or *polarimetry*. Hence, polarimetric instruments inevitably require the information about the polarization properties of the optical system or the *instrumental polarization*. This can be determined through experiments, analysis or a combination of both. As the telescopes become larger and instruments become more complex, the instrumental polarization analysis has become all the more crucial (e.g., [Anche et al., 2015](#); [Harrington & Sueoka, 2016](#)).

## Summary

The Python package `PyAstroPol` provides a means to analyze the polarization properties of a given optical system with relative ease and minimal dependencies. The simple end goal is to calculate the Mueller matrix (e.g., [Gil & Ossikovski, 2016](#)) of the optical system. Data pipelines of the astronomical instruments can easily distribute this package alongside their polarimetric calibration routines.

In the polarization analysis of the astronomical telescopes, various approaches have been adopted depending on the complexity of the system. A significant part of the complexity is because the instrumental polarization is often time-dependent. For the solar telescopes with Coelostats, Mueller matrices were analytically derived as a function of time (e.g., [Balasubramaniam et al., 1985](#); [Beck et al., 2005](#)). For Thirty Meter Telescope (TMT), a combination of analytical and numerical methods is used ([Anche et al., 2015](#)). However, for Daniel K. Inouye Solar Telescope (DKIST), Zemax – a commercial software – is used, along with the team’s in-house tools ([Harrington & Sueoka, 2016](#)).

`PyAstroPol` aims to combine the better parts of the aforementioned programs: it is an open-source tool written in the Python programming language and offers a range of functions to model the astronomical optics. There is no open-source software which has polarization ray propagation features as per my knowledge. Hence, Zemax OpticStudio® modelling is used for comparison. The salient features and important limitations are listed below.

Salient features :

- `PyAstroPol` calculates the Mueller matrix of a given imaging-type optical system, by coherently adding the electric field vectors after propagation.
- Astronomical sources can be directly placed in the model using relevant coordinates, namely, declination, hour angle and latitude of the telescope site.

- Off-axis components are included, as they have a significant effect on polarization.
- Effect of multi-layered coatings, such as oxide layers and protective coatings, on the state of polarization is included. These are also significant in the polarization analysis of the astronomical optics (e.g., [Harten et al., 2009](#)).
- All the data, such as points of incidence, polarization directions, complex electric field values and more, are readily available to the user for any further analysis.
- Material refractive index information can be downloaded from the popular online source <https://refractiveindex.info/> as .csv. It can be formatted and used with this software.
- Spot diagram is possible at any instance of the beam path, as a by-product of ray tracing.

#### Important Limitations:

- All the analysis uses strictly the ray treatment. Hence, all the limitations of the rays optics shall be applicable.
- Only circular optics (apertures) can be modelled at the moment.
- Birefringent components are not included yet. The justification for this choice is that the behaviour of the birefringent components is fairly straightforward as they strongly polarize the light.
- The visualization features are limited.
- PyAstroPol is neither a design software nor interactive. That is, the user must know the optical system that is to be analyzed, and the system must be updated every time a component is changed.

Results of the code have been verified against previous works ([Pruthvi et al., 2018](#)), and Zemax OpticStudio®. A variety of examples have been provided, and they should facilitate the quick-start. One of the examples also provides the aforementioned comparison with commercial software.

## Acknowledgements

This work has been carried out as a part of the ongoing project *Jets in the solar atmosphere*, funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Projektnummer 407727365. I thank DVS Phanindra (IIA, Bengaluru) and V. Sreekanth Reddy (CHESS, Hyderabad) for the discussion, and Mathias Waidele (KIS, Freiburg) for his feedback. I also thank the PI of the DFG project Markus Roth (KIS, Freiburg) for the opportunity to carry out this work. I thank reviewers [@caldarolamartin](#), [@aquilesC](#) and [@mwrcraig](#), and editor [@pibion](#) for their valuable feedback in improving this package.

## References

- Anche, R. M., Anupama, G. C., Reddy, K., Sen, A., Sankarasubramanian, K., Ramaprakash, A. N., Sengupta, S., Skidmore, W., Atwood, J., Tirupathi, S., & Pandey, S. B. (2015). Analytical modelling of Thirty Meter Telescope optics polarization. In K. Bhattacharya (Ed.), *International conference on optics and photonics 2015* (Vol. 9654, pp. 45–52). International Society for Optics; Photonics; SPIE. <https://doi.org/10.1117/12.2181397>
- Balasubramaniam, K. S., Venkatakrishnan, P., & Bhattacharyya, J. C. (1985). Measurement of vector magnetic fields. I - Theoretical approach to the instrumental polarisation of the Kodaikanal Solar Tower. *99*, 333–348. <https://doi.org/10.1007/BF00157317>

- Beck, C., Schlichenmaier, R., Collados, M., Bellot Rubio, L., & Kentischer, T. (2005). A polarization model for the German Vacuum Tower Telescope from in situ and laboratory measurements. *Astronomy & Astrophysics*, 443, 1047–1053. <https://doi.org/10.1051/0004-6361:20052935>
- Gil, J. J., & Ossikovski, R. (2016). *Polarized Light and the Mueller Matrix Approach*. CRC Press. <https://doi.org/10.1201/b19711>
- Harrington, D., & Sueoka, S. R. (2016). *Polarization modeling and predictions for DKIST part 1: Telescope and example instrument configurations* (Vol. 9912, p. 99126U). Proc SPIE. <https://doi.org/10.1117/12.2233176>
- Harten, G. van, Snik, F., & Keller, C. U. (2009). Polarization Properties of Real Aluminum Mirrors, I. Influence of the Aluminum Oxide Layer. *Publications of the Astronomical Society of the Pacific*, 121(878), 377–383. <https://doi.org/10.1086/599043>
- Pruthvi, H., Krishnappa, N., Belur, R., & Kadagattor, R. B. (2018). Solar spectropolarimetry of Ca II 8542 Å line: polarimeter development, calibration, and preliminary observations. *Journal of Astronomical Telescopes, Instruments, and Systems*, 4(04), 1. <https://doi.org/10.1117/1.JATIS.4.4.048002>