

# hawen: time-harmonic wave modeling and inversion using hybridizable discontinuous Galerkin discretization

Florian Faucher<sup>1</sup>

<sup>1</sup> Faculty of Mathematics, University of Vienna, Oskar-Morgenstern-Platz 1, A-1090 Vienna, Austria.

DOI: [10.21105/joss.02699](https://doi.org/10.21105/joss.02699)

**Software**

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

**Editor:** [Marie E. Rognes](#) ↗

**Reviewers:**

- [@chennachaos](#)
- [@yangbai90](#)

**Submitted:** 01 September 2020


**Published:** 24 January 2021

**License**

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

## Summary

Many applications such as seismic and medical imaging, material sciences, or helioseismology and planetary science, aim to reconstruct properties of a non directly accessible or non-visible interior. For this purpose, they rely on waves whose propagation through a medium interrelates with the physical properties (density, sound speed, etc.) of this medium. The methodology for imaging with waves comprises of two main stages illustrated in [Figure 1](#). In the data acquisition stage ([Figure 1a](#)), the medium response to probing waves is recorded (e.g., seismic waves from Earthquakes recorded by ground network). In the second stage, we rely on a reconstruction procedure which iteratively updates an initial model of physical parameters, so that numerical simulations approach the measurements ([Figure 1b](#)). This procedure is employed, for instance, for seismic (reconstruction of subsurface layers) and medical (disease diagnostic) imaging, see the introduction of [Faucher & Scherzer \(2020\)](#) and the references therein.



**Figure 1:** Illustration of inverse wave problems. **a)** Acquisition stage: probing waves are sent though the medium, and the medium’s response is recorded by devices positioned on a portion of the domain. **b)** The reconstruction algorithm starts from initial properties and compares simulations of wave propagation with the measurements, then iteratively updates those properties. The green block corresponds to the forward problem (modeling the propagation of waves) and the orange ones to the iterative correction. *hawen* solves both the forward and inverse problems associated with time-harmonic waves.

## Statement of need

`hawen` is designed to address large-scale inverse wave problems, and includes

1. simulating time-harmonic waves in heterogeneous media (e.g., visco-acoustic and visco-elastic propagation, modal equations in helioseismology);
2. performing iterative minimization to solve the inverse problem, cf. [Figure 1](#).


It combines MPI and OpenMP parallelism, it is deployed on supercomputers and has been used in studies on seismic imaging ([Faucher, Alessandrini, et al., 2020](#); [Faucher, De Hoop, et al., 2020](#); [Faucher & Scherzer, 2020](#)), as well as in helioseismology ([Barucq et al., 2019, 2020](#); [Barucq, Faucher, Fournier, et al., 2020a, 2020b](#)). The software works with an input *parameter file*: a text file listing the problem configuration (dimension, choice of viscous model, etc.), such that the computational details are transparent for the users. `hawen` has its dedicated [User's guide](#) ([Faucher, 2020](#)) that details its utilization and all the available functionalities, it is available on the software dedicated [website](#), which contains illustrations, installation guide and tutorials.

One specificity of `hawen` is to implement the hybridizable discontinuous Galerkin (HDG) method ([Arnold et al., 2002](#); [Cockburn et al., 2009](#)) for the discretization of the wave equations. It helps reduce the cost of the computations by providing smaller linear systems compared to, e.g., finite elements, depending on the order of approximation ([Faucher & Scherzer, 2020](#); [Kirby et al., 2012](#)). For seismic applications, current software mostly relies on the spectral element method ([Komatitsch & Tromp, 1999](#); [Komatitsch & Vilotte, 1998](#)), such as `specfem`, or on finite differences discretization.

The software handles unstructured domains (allowing input meshes of different formats) and thus accounts for complex geometry. In addition, `hawen` relies on the massively parallel sparse direct solver MUMPS ([Amestoy et al., 2001, 2006](#)) for the matrix factorization, hence handles problems with many right-hand sides. Despite the technicality of the underlying methods, the purpose of `hawen` is to provide a unified and evolutive framework to address large-scale visco-elastic and visco-acoustic inverse problems, with a user-friendly interface using the input parameter files.

## Modeling the propagation of waves

The first feature of `hawen` is to simulate the propagation of time-harmonic waves in different types of medium: this is the forward problem in [Figure 1](#). The propagation of waves is characterized by a Partial Differential Equation which depends on the type of the waves (e.g., mechanical or electromagnetic), and on the type of medium considered (e.g., fluid or solid) ([Carcione, 2007](#); [Faucher, 2017](#); [Slawinski, 2010](#)). `hawen` handles visco-acoustic and visco-elastic propagation, as well as waves propagation for helioseismology, these are described in the [User's documentation](#) ([Faucher, 2020](#)). Our implementation is based upon the HDG method and the computational steps are illustrated in [Figure 2](#).



**Figure 2:** Illustration of the computational steps for the numerical resolution of the forward problem. For simplicity, we illustrate with an homogeneous medium.

## Inverse problem via iterative minimization

In the forward problem, the objective is to simulate the propagation of waves, given the medium physical properties. Conversely, the objective of the inverse problem is to recover the physical properties, given some observations of waves, see [Figure 1](#). To address the inverse problem, `hawn` solves the minimization problem:

$$\min_m \mathcal{J}(m) \quad \text{with} \quad \mathcal{J}(m) = \text{dist}(\mathcal{F}(m), d). \quad (1)$$

The misfit function  $\mathcal{J}$  is defined to evaluate a distance function comparing the observed data  $d$  with simulations of wave propagation restricted to the position of the measurements:  $\mathcal{F}(m)$ , where  $m$  represents the physical properties. The iterative minimization scheme successively updates the physical properties used for the simulations ([Figure 1b](#)), and follows a Newton-type method ([Nocedal & Wright, 2006](#)). In the context of seismic imaging, it is the Full Waveform Inversion method, cf. [Virieux & Operto \(2009\)](#).

`hawn` offers several options to conduct the iterative minimization, such as the choice of misfit function and method to conduct the minimization. These are further listed and detailed in the [software documentation](#) ([Faucher, 2020](#)).

## Acknowledgements

The research of FF is supported by the Austrian Science Fund (FWF) under the Lise Meitner fellowship M 2791-N. We acknowledge the use of the [Vienna Scientific Clusters vsc3 and vsc4](#) for the numerical applications, as well as with the TGCC cluster via the GENCI resource allocation project AP010411013.

## References

- Amestoy, P. R., Duff, I. S., L'Excellent, J.-Y., & Koster, J. (2001). A fully asynchronous multifrontal solver using distributed dynamic scheduling. *SIAM Journal on Matrix Analysis and Applications*, 23(1, 1), 15–41. <https://doi.org/10.1137/S0895479899358194>
- Amestoy, P. R., Guermouche, A., L'Excellent, J.-Y., & Pralet, S. (2006). Hybrid scheduling for the parallel solution of linear systems. *Parallel Computing*, 32(2, 2), 136–156. <https://doi.org/10.1016/j.parco.2005.07.004>
- Arnold, D. N., Brezzi, F., Cockburn, B., & Marini, L. D. (2002). Unified analysis of discontinuous Galerkin methods for elliptic problems. *SIAM Journal on Numerical Analysis*, 39(5), 1749–1779. <https://doi.org/10.1137/S0036142901384162>
- Barucq, H., Faucher, F., Fournier, D., Gizon, L., & Pham, H. (2020a). Efficient and accurate algorithm for the full modal Green's kernel of the scalar wave equation in helioseismology. *SIAM Journal on Applied Mathematics*, 80, 2657–2683. <https://doi.org/10.1137/20M1336709>
- Barucq, H., Faucher, F., Fournier, D., Gizon, L., & Pham, H. (2020b). *Efficient computation of the modal outgoing Green's kernel for the scalar wave equation in helioseismology* (Research Report RR-9338; pp. 1–85). Inria Bordeaux Sud-Ouest; Magique 3D; Max-Planck Institute for Solar System Research. <https://hal.archives-ouvertes.fr/hal-02544701>
- Barucq, H., Faucher, F., & Pham, H. (2019). *Outgoing solutions to the scalar wave equation in helioseismology* (Research Report RR-9280; pp. 1–136). Inria Bordeaux Sud-Ouest ; Project-Team Magique3D. <https://hal.archives-ouvertes.fr/hal-02168467>
- Barucq, H., Faucher, F., & Pham, H. (2020). Outgoing solutions and radiation boundary conditions for the ideal atmospheric scalar wave equation in helioseismology. *ESAIM: Mathematical Modelling and Numerical Analysis*, 54, 1111–1138. <https://doi.org/10.1051/m2an/2019088>
- Carcione, J. M. (2007). *Wave fields in real media: Wave propagation in anisotropic, anelastic, porous and electromagnetic media*. Elsevier.
- Cockburn, B., Gopalakrishnan, J., & Lazarov, R. (2009). Unified hybridization of discontinuous galerkin, mixed, and continuous galerkin methods for second order elliptic problems. *SIAM Journal on Numerical Analysis*, 47(2), 1319–1365. <https://doi.org/10.1137/070706616>
- Faucher, F. (2017). *Contributions to seismic full waveform inversion for time harmonic wave equations: Stability estimates, convergence analysis, numerical experiments involving large scale optimization algorithms* (pp. 1–400) [PhD thesis, Université de Pau et Pays de l'Ardour]. <https://hal.archives-ouvertes.fr/tel-01807861>
- Faucher, F. (2020). *Hawen (time-HARmonic wave modELing and INversion using hybridizable discontinuous galerkin discretization) user's guide* (Version 0.1.0) [Computer software]. [https://ffaucher.gitlab.io/hawen-website/download/hawen\\_users-guide\\_documentation.pdf](https://ffaucher.gitlab.io/hawen-website/download/hawen_users-guide_documentation.pdf)
- Faucher, F., Alessandrini, G., Barucq, H., Hoop, M. de, Gaburro, R., & Sincich, E. (2020). Full reciprocity-gap waveform inversion, enabling sparse-source acquisition. *Geophysics*, 85(6), 1–82. <https://doi.org/10.1190/geo2019-0527.1>
- Faucher, F., De Hoop, M. V., & Scherzer, O. (2020). Reciprocity-gap misfit functional for distributed acoustic sensing, combining data from passive and active sources. *Geophysics*, 86, 1–46. <https://doi.org/10.1190/geo2020-0305.1>
- Faucher, F., & Scherzer, O. (2020). Adjoint-state method for Hybridizable Discontinuous Galerkin discretization: Application to the inverse acoustic wave problem. *Computer*

- Methods in Applied Mechanics and Engineering*, 372, 113406. <https://doi.org/10.1016/j.cma.2020.113406>
- Kirby, R. M., Sherwin, S. J., & Cockburn, B. (2012). To CG or to HDG: A comparative study. *Journal of Scientific Computing*, 51(1), 183–212. <https://doi.org/10.1007/s10915-011-9501-7>
- Komatitsch, D., & Tromp, J. (1999). Introduction to the spectral element method for three-dimensional seismic wave propagation. *Geophysical Journal International*, 139(3), 806–822. <https://doi.org/10.1046/j.1365-246x.1999.00967.x>
- Komatitsch, D., & Vilotte, J.-P. (1998). The spectral element method: An efficient tool to simulate the seismic response of 2D and 3D geological structures. *Bulletin of the Seismological Society of America*, 88(2), 368–392.
- Nocedal, J., & Wright, S. J. (2006). *Numerical optimization* (2nd ed.). Springer Series in Operations Research. <https://doi.org/10.1007/b98874>
- Slawinski, M. A. (2010). *Waves and rays in elastic continua*. World Scientific. <https://doi.org/10.1142/7486>
- Virieux, J., & Operto, S. (2009). An overview of full-waveform inversion in exploration geophysics. *Geophysics*, 74(6), WCC1–WCC26. <https://doi.org/10.1190/1.3238367>