

Qlunc: Quantification of lidar uncertainty

Francisco Costa¹, Andrew Clifton¹, Nikola Vasiljevic², and Ines Würth¹

DOI: [10.21105/joss.03211](https://doi.org/10.21105/joss.03211)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Daniel S. Katz](#) ↗

Reviewers:

- [@adi3](#)
- [@antviro](#)

Submitted: 12 March 2021

Published: 28 October 2021

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

¹ Stuttgart Wind Energy (SWE), Allmandring 5b, 70569 Stuttgart, Germany ² DTU Wind Energy, Frederiksborgvej 399, 4000 Roskilde Denmark

Summary

Wind lidar is a flexible and versatile remote sensing device for wind energy applications (Beck & Kühn, 2017) that measures the wind vector remotely using laser light backscattered from aerosols. It is a key tool for wind energy and meteorology. As with any measurement method, it is essential to estimate its uncertainty. Qlunc, which stands for **Q**uantification of lidar **u**ncertainty, is an open-source Python-based tool to create a digital twin of lidar hardware, and to estimate the uncertainty of wind lidar wind speed measurements.

Qlunc contains models of the uncertainty contributed by individual lidar components and modules (represented by Python objects, which in turn represent physical lidar objects), that then are combined, considering their different natures, to estimate the uncertainties in wind lidar measurements. The modules are based on the OpenLidar architecture (Clifton et al., 2019) and can be easily adapted for particular use cases thanks to the modularity of the code (see Figure 1). The terminology for the components and modules defined within Qlunc has also been aligned with a community-driven wind lidar ontology, which is in development (N. Vasiljevic, 2021a, 2021b).




Figure 1: Qlunc basic structure.

The first release is focused on velocity azimuth display (VAD)(Browning & Wexler, 1968) scans and forward-looking nacelle-mounted measuring modes, which are common wind-energy-industry applications. Besides uncertainty estimations, Qlunc’s functions could be extended for other applications, for example to compare different wind velocity vector calculation methods. This, combined with the underlying open-source code, defines an attractive scenario for sharing knowledge and fostering collaboration on wind lidars.

Statement of Need

Wind lidars are measuring devices, and as for any other measuring systems, their measurements have uncertainties (Borraccino & Courtney, 2016). Therefore, as already stated, it is crucial to assess their measurement uncertainty in order to increase confidence in lidar technology.

Measurement uncertainty means doubt about the validity of the result of a measurement (JCGM, 2008). It represents the dispersion of the values attributed to a measurand. The ability to simulate uncertainty through a model such as Qlunc is important for judging measurement data but can also be useful for designing and setting up experiments and optimizing lidar design. Because wind lidar is important for wind energy applications (Clifton et al., 2018), better models for wind lidar hardware (e.g., Qlunc) and measurement processes (e.g., through MOCALUM (Nikola Vasiljevic, 2020) or YADDUM (Nikola Vasiljevic, 2019), with which Qlunc can feasibly combine) will directly contribute to the adoption of wind lidar for wind energy applications.

This project is influenced by fundamental open science principles (Bot et al., 2018). The scope is to create an open, standardized and collaborative framework to describe both generic and specific lidar architectures, characterize lidar uncertainties, and provide the tools for others to contribute within this framework.

Future development roadmap

Over the next year, we plan to implement further lidar hardware modules in the model and compute their combined uncertainties. In addition, we will identify main data processing methods and include those that we consider the highest contributors to uncertainty.

We also plan to further align the terminology used in Qlunc with the IEA Wind Task 32 controlled vocabulary for wind lidar (N. Vasiljevic et al., 2021). This will make it easier for users to understand what each of the modules and components do, and promotes interoperability.

All documentation from the project, tutorials, and raw code will be published through a website, to enable users to dive into the numerical framework and get used to the Qlunc routines. We welcome contributions from the wind lidar community.

Acknowledgements

This work is part of the LIKE (Lidar Knowledge Europe) project. The project LIKE H2020-MSCA-ITN-2019, Grant number 858358 is funded by the European Union.

References

- Beck, H., & Kühn, M. (2017). Dynamic Data Filtering of Long-Range Doppler LiDAR Wind Speed Measurements. *Remote Sensing*, 9(6). <https://doi.org/10.3390/rs9060561>
- Borraccino, A., & Courtney, M. (2016). *Calibration report for ZephIR Dual Mode lidar (unit 351)*. <https://doi.org/10.13140/RG.2.1.1658.2005>
- Bot, G., Heller, L., datawomanHUB, mcancellieri, Kramer, B., Ross-Hellauer, T., ilabastida, helenebr, Fernandes, P., & Tennant, J. (2018). *Open Science Training Handbook (Version 1.1)* [Computer software]. Zenodo. <https://doi.org/10.5281/zenodo.1212538>
- Browning, K. A., & Wexler, R. (1968). The Determination of Kinematic Properties of a Wind Field Using Doppler Radar. *Journal of Applied Meteorology and Climatology*. [https://doi.org/10.1175/1520-0450\(1968\)007%3C0105:TDOKPO%3E2.0.CO;2](https://doi.org/10.1175/1520-0450(1968)007%3C0105:TDOKPO%3E2.0.CO;2)

- Clifton, A., Clive, P., Gottschall, J., Schlipf, D., Simley, E., Simmons, L., Stein, D., Trabucchi, D., Vasiljevic, N., & Würth, I. (2018). IEA Wind Task 32: Wind Lidar Identifying and Mitigating Barriers to the Adoption of Wind Lidar. *Remote Sensing*, 10(3). <https://doi.org/10.3390/rs10030406>
- Clifton, A., Vasiljevic, N., Wuerth, I., Raach, S., Haizmann, F., & Fuerst, H. (2019). *The OpenLidar Initiative for collaboration on wind lidar hardware and software*. <https://doi.org/10.5281/zenodo.3414197>
- JCGM. (2008). *Evaluation of measurement data: Guide to the expression of uncertainty in measurement*. International Organization for Standardization. ISBN: [9267101889](https://doi.org/10.1002/978-3-7089-1898-9)
- Vasiljevic, N. (2021a). *sheet2rdf: automatic workflow for generation of RDF vocabularies from Google sheets* (Version v0.1) [Computer software]. Zenodo. <https://doi.org/10.5281/zenodo.4432136>
- Vasiljevic, Nikola. (2020). *mocalum: A Python package for Monte-Carlo lidar uncertainty modeling*. Zenodo. <https://doi.org/10.5281/zenodo.3823878>
- Vasiljevic, Nikola. (2019). *YADDUM* (Version v0.2.0) [Computer software]. Zenodo. <https://doi.org/10.5281/zenodo.3580749>
- Vasiljevic, N. (2021b). *OntoStack*. <http://data.windenergy.dtu.dk/ontologies/view/en/>
- Vasiljevic, N., Clifton, A., & Costa, F. (2021). *IEA Wind Task 32 Wind Lidar Ontology*. <http://vocab.ieawindtask32.org/ontolidar/en/>