

MMM: A C++ library for simulating large herbivores

Traylor, Wolfgang¹

¹ Senckenberg Biodiversity and Climate Research Centre, Frankfurt am Main, Germany

DOI: [10.21105/joss.03631](https://doi.org/10.21105/joss.03631)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Patrick Diehl](#) ↗

Reviewers:

- [@dirmeier](#)
- [@dilawar](#)

Submitted: 02 July 2021

Published: 15 December 2021

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

The Modular Megafauna Model (MMM) simulates populations of large, terrestrial herbivores (megafauna) through space and time.

Herbivore cohorts feed, grow, reproduce, and die in daily simulation cycles. Based on these physiological processes herbivore numbers rise and fall dynamically. The model thus does not prescribe carrying capacity, but instead simulates herbivore densities in a mechanistic, bottom-up approach.

Statement of Need

Mechanistic modeling can help us understand the drivers behind real-world population dynamics. Bottom-up models can point out which physiological processes are understudied and which mechanisms appear most important for emerging population effects ([DeAngelis & Mooij, 2003](#)). Once a model is sufficiently mature, its predictions can inform nature conservation management—an application that could prove useful in light of ongoing global defaunation ([Dirzo et al., 2014](#)).

The currently implemented model concepts originate in large parts from [Pachzelt et al. \(2013\)](#) and the earlier works by [Illius & O'Connor \(2000\)](#) and [Illius & Gordon \(1999\)](#). [Pachzelt et al. \(2013\)](#) integrated African grazers into LPJ-GUESS, a dynamic global vegetation model (DGVM) ([Benjamin Smith et al., 2001](#)). Later studies have implemented conceptually similar grazer models for other DGVMs: [Dangal et al. \(2017\)](#) for DLEM and [Zhu et al. \(2018\)](#) for ORCHIDEE. However, to my knowledge, none of these implementations are reusable across different vegetation models.

The author is currently using MMM integrated with LPJ-GUESS ([Benjamin Smith et al., 2001](#); [B. Smith et al., 2014](#)), using daily grass growth ([Boke-Olén et al., 2018](#)).¹ Here, LPJ-GUESS simulates grass as forage for herbivores. This particular project is concerned with simulating potential densities of mammoths, steppe bison, and horse in the last ice age. However, MMM could also be parametrized for extant herbivores, such as cattle, Scandinavian reindeer, or Mongolian gazelles. MMM can also be coupled with dynamic vegetation models other than LPJ-GUESS.

Features

MMM is a C++ library meant to be coupled with a dynamic vegetation model into a complete ecosystem model. Currently, the only forage is grass. The vegetation model simulates

¹The proprietary LPJ-GUESS source code, including the MMM-coupled version, may be available upon request at Lund University (<https://iis4.nateko.lu.se/lpj-guess/>).

the amount of available forage, which the herbivores then consume. In addition, the vegetation model provides information about environmental conditions, such as air temperature. This way, herbivores and vegetation dynamically influence each other, namely through forage removal and nutrient cycling. MMM ships with a very simple vegetation model that demonstrates how the coupling can be implemented.

Herbivores are simulated in distinct spatial units, which have no absolute area size because all calculations are done on a per-area basis. That means that MMM itself is not spatially explicit and makes no assumptions about the actual size of the area inhabited by herbivores. It is up to the vegetation model to give these spatial units meaning by linking them to spatially explicit entities, such as grid cells. With this flexibility, MMM can be used for studies on different scales, from local to continental.

Modularity is a primary design goal of the library. Through the instruction file², users can turn mechanisms on or off and parametrize herbivore species or herbivore functional types. Parameters include, for example, body mass, components of energy expenditure, maximum feed intake, background mortality, body fat reserves, etc. There are no hard-coded parameters; all can be defined in the instruction file. MMM's flexible framework allows developers to integrate new mechanisms, such as a more detailed energy budget model, mortality from hunting and predation, or a new forage type.

While monolithic ecosystem models can easily become “black boxes,” whose internal mechanisms have grown too complex to be understood intuitively, a modular model is more transparent. Developing mechanistic ecosystem models is typically an exploratory, iterative process. For a specific study, the modeler has to adjust parameters and mechanisms of a given model, either manually or programmatically. In this process it is crucial that the modeler can increase model complexity step by step, just enough to represent the mechanisms important for the research question.

Thanks to its modular design and its stable library interface, MMM can stay backward-compatible in future versions, and the same codebase can be used for different studies and different vegetation models. This benefits reproducibility in two ways. First, after bugs have been fixed, previous analyses can easily be reexecuted. Second, simulations can be repeated with other vegetation models in order to understand how their different assumptions impact plant–herbivore dynamics.

Acknowledgements

I thank my PhD supervisor Thomas Hickler and my colleagues Adrian Pachzelt, Matthew Forrest, and Theresa Stratmann for their support in model development and implementation.

References

- Boke-Olén, N., Lehsten, V., Abdi, A. M., Ardö, J., & Khatir, A. A. (2018). Estimating grazing potentials in Sudan using daily carbon allocation in dynamic vegetation model. *Rangeland Ecology & Management*, 71(6), 792–797. <https://doi.org/10.1016/j.rama.2018.06.006>
- Dangal, S. R. S., Tian, H., Lu, C., Ren, W., Pan, S., Yang, J., Di Cosmo, N., & Hessel, A. (2017). Integrating herbivore population dynamics into a global land biosphere model: Plugging animals into the earth system. *Journal of Advances in Modeling Earth Systems*, 9(8), 2920–2945. <https://doi.org/10.1002/2016MS000904>

²Example instruction files are included in the code repository: https://github.com/wtraylor/modular_mega fauna_model/tree/master/examples

- DeAngelis, D. L., & Mooij, W. M. (2003). In praise of mechanistically rich models. In C. D. Canham, J. J. Cole, & W. K. Lauenroth (Eds.), *Models in ecosystem science* (pp. 62–82). Princeton University Press. <https://doi.org/10.2307/j.ctv1dwq0tq.9>
- Dirzo, R., Young, H. S., Galetti, M., Ceballos, G., Isaac, N. J. B., & Collen, B. (2014). Defaunation in the Anthropocene. *Science*, *345*(6195), 401–406. <https://doi.org/10.1126/science.1251817>
- Illius, A. W., & Gordon, I. J. (1999). Scaling up from functional response to numerical response in vertebrate herbivores. In H. Olf, V. K. Brown, & R. H. Drent (Eds.), *Herbivores: Between plants and predators: The 38th symposium of the British Ecological Society 1997* (pp. 397–425). British Ecological Society; Blackwell Science.
- Illius, A. W., & O'Connor, T. G. (2000). Resource heterogeneity and ungulate population dynamics. *Oikos*, *89*(2), 283–294. <https://doi.org/10.1034/j.1600-0706.2000.890209.x>
- Pachzelt, A., Rammig, A., Higgins, S., & Hickler, T. (2013). Coupling a physiological grazer population model with a generalized model for vegetation dynamics. *Ecological Modelling*, *263*, 92–102. <https://doi.org/10.1016/j.ecolmodel.2013.04.025>
- Smith, Benjamin, Prentice, I. C., & Sykes, M. T. (2001). Representation of vegetation dynamics in the modelling of terrestrial ecosystems: Comparing two contrasting approaches within European climate space. *Global Ecology and Biogeography*, *10*(6), 621–637. <https://doi.org/10.1046/j.1466-822X.2001.t01-1-00256.x>
- Smith, B., Wårlind, D., Arneeth, A., Hickler, T., Leadley, P., Siltberg, J., & Zaehle, S. (2014). Implications of incorporating N cycling and N limitations on primary production in an individual-based dynamic vegetation model. *Biogeosciences*, *11*(7), 2027–2054. <https://doi.org/10.5194/bg-11-2027-2014>
- Zhu, D., Ciais, P., Chang, J., Krinner, G., Peng, S., Viovy, N., Peñuelas, J., & Zimov, S. (2018). The large mean body size of mammalian herbivores explains the productivity paradox during the last glacial maximum. *Nature Ecology & Evolution*, *2*(4), 640–649. <https://doi.org/10.1038/s41559-018-0481-y>