




# HostPhot: global and local photometry of galaxies hosting supernovae or other transients

Tomás E. Müller-Bravo <sup>1,2</sup> and Lluís Galbany <sup>1,2</sup>

1 Institute of Space Sciences (ICE, CSIC), Campus UAB, Carrer de Can Magrans, s/n, E-08193 Barcelona, Spain 2 Institut d'Estudis Espacials de Catalunya (IEEC), E-08034 Barcelona, Spain

DOI: [10.21105/joss.04508](https://doi.org/10.21105/joss.04508)

## Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

---

Editor: [Arfon Smith](#) 

## Reviewers:

- [@Onoddil](#)
- [@krachyon](#)

Submitted: 31 May 2022

Published: 15 August 2022

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## Summary

Type Ia supernovae (SNe Ia) have assumed a fundamental role as cosmological distance indicators since the discovery of the accelerating expansion rate of the universe ([Perlmutter et al., 1999](#); [Riess et al., 1998](#)). Correlations between their optical peak luminosity, the decline rate of their light curves and their optical colours allow them to be standardised, reducing their observed r.m.s scatter (e.g. [Phillips, 1993](#); [Tripp, 1998](#)). Over a decade ago, the optical peak luminosity of SNe Ia was found to correlate with host galaxy stellar mass, further improving their standardisation ([Kelly & others, 2010](#); [Lampeitl et al., 2010](#); [Sullivan et al., 2010](#)). Since then, host galaxy properties have been used in cosmological analyses of SNe Ia ([Betoule et al., 2014](#); [Brout et al., 2019](#); [Scolnic et al., 2018](#)) and tremendous effort has gone into finding the property, such as star formation rate ([Rigault et al., 2013](#)), that fundamentally drives the correlation between SNe Ia and their host galaxies. Furthermore, it has been noted that the local environment in which the progenitors of SNe Ia evolve is much better at reducing the scatter in estimated distances than the global environment, i.e., the whole galaxy ([Kelsey et al., 2021](#); [Roman et al., 2018](#)). Therefore, the study of the effect of environment on SNe Ia is an active field of research and key in future cosmological analyses.

## Statement of need

HostPhot is an open-source Python package for measuring galaxy photometry, both locally and globally. Galaxy photometry is fundamental as it is commonly used to estimate the galaxy parameters, such as stellar mass and star formation rate. However, the codes used to calculate photometry by different groups can vary and there is no dedicated package for this. The API for HostPhot allows the user to extract public image cutouts of surveys, such as the Panoramic Survey Telescope and Rapid Response System (Pan-STARRS) Data Release 1 (PS1), Dark Energy Survey (DES) and Sloan Digital Sky Survey (SDSS). Different sets of filters are available depending on the chosen survey: *grizy* for PS1, *grizY* for DES and *ugriz* for SDSS. All photometry is corrected for Milky Way dust extinction. Furthermore, HostPhot also works with private data obtained by the user and can even be easily modified to include other surveys.

The major novelty of HostPhot is dealing with low-redshift galaxies ( $z < 0.1$ ) as obtaining photometry of these is not as simple as those at higher redshift. Foreground stars can be in the line of sight of nearby galaxies, making the extraction of the photometry a complex procedure. In addition, low-redshift galaxies have visible structures, while at high redshift they just look like simple ellipses. HostPhot is able to detect sources in the images, cross-match them with catalogs of stars (e.g., Gaia ([Gaia Collaboration, 2016](#))) and remove them by applying a convolution with a 2D Gaussian kernel. This process ensures that only stars (and in some cases other galaxies that are not of interest) are removed, keeping the structure of the galaxy intact.

HostPhot can calculate the photometry of an entire galaxy (global) or in a given circular aperture (local) and it heavily relies on the Astropy (Astropy Collaboration et al., 2018, 2013) and Photutils (Bradley et al., 2021) packages for this. Local photometry can be calculated for different circular apertures in physical units (e.g., 4 kpc) at the redshift of the given object. In addition, as the physical size depends on the assumed cosmology, the cosmological model can be changed by the user, suiting their needs. On the other hand, for the global photometry, the user can choose between using a different aperture for each filter/image or a common aperture for all the filters/images. For the latter, HostPhot coadds images in the desired filters, as selected by the user (e.g., *riz*), and estimates the common aperture parameters from the coadd image. The aperture used for the global photometry can also be optimised, by increasing the size until the change in flux is negligible, encompassing the entire galaxy. In a few cases, nearby galaxies can have very complex structures. HostPhot offers the option of interactively setting the aperture via an intuitive GUI. This option also allows the user to test how the change in aperture shape can affect the calculated photometry.

HostPhot is user-friendly and well documented<sup>1</sup>, which allows the community to easily contribute to this package. HostPhot is already being used by different groups, such as HostFlows<sup>2</sup> and DES, and will allow the supernova community to find exciting new scientific discoveries with future cosmological analyses. Finally, although HostPhot is mainly aimed at supernova science, it can be used in other fields in astronomy as well.

Apart from Astropy and Photutils (Bradley et al., 2021), HostPhot also relies on sep (Barbary, 2016b) for global photometry, Astroquery (Ginsburg et al., 2019) for image downloading and cross-matching with catalogs, reproject<sup>3</sup> for the coadds, extinction (Barbary, 2016a) and sfdmap<sup>4</sup> for extinction correction. Finally, HostPhot makes use of the following packages as well: numpy (Harris et al., 2020), matplotlib (Hunter, 2007), pandas (McKinney & others, 2010), pyvo (Graham et al., 2014), ipywidgets<sup>5</sup> and ipympl<sup>6</sup>.

## Acknowledgements

TEMB and LG acknowledge financial support from the Spanish Ministerio de Ciencia e Innovación (MCIN), the Agencia Estatal de Investigación (AEI) 10.13039/501100011033 under the PID2020-115253GA-I00 HOSTFLOWS project, and from Centro Superior de Investigaciones Científicas (CSIC) under the PIE project 20215AT016, and the I-LINK 2021 LINKA20409. TEMB and LG are also partially supported by the program Unidad de Excelencia María de Maeztu CEX2020-001058-M. LG also acknowledges MCIN, AEI and the European Social Fund (ESF) “Investing in your future” under the 2019 Ramón y Cajal program RYC2019-027683-I.

## References

- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., Günther, H. M., Lim, P. L., Crawford, S. M., Conseil, S., Shupe, D. L., Craig, M. W., Dencheva, N., Ginsburg, A., VanderPlas, J. T., Bradley, L. D., Pérez-Suárez, D., de Val-Borro, M., Aldcroft, T. L., Cruz, K. L., Robitaille, T. P., Tollerud, E. J., ... Astropy Contributors. (2018). The Astropy Project: Building an Open-science Project and Status of the v2.0 Core Package. *The Astronomical Journal*, 156(3), 123. <https://doi.org/10.3847/1538-3881/aabc4f>
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., Greenfield, P., Droettboom, M., Bray, E., Aldcroft, T., Davis, M., Ginsburg, A., Price-Whelan, A. M., Kerzendorf, W. E., Conley,

<sup>1</sup><https://hostphot.readthedocs.io/en/latest/>

<sup>2</sup><https://hostflows.github.io/>

<sup>3</sup><https://pypi.org/project/reproject/>

<sup>4</sup><https://github.com/kbarbary/sfdmap>

<sup>5</sup><https://github.com/jupyter-widgets/ipywidgets>

<sup>6</sup><https://github.com/matplotlib/ipympl>

- A., Crighton, N., Barbary, K., Muna, D., Ferguson, H., Grollier, F., Parikh, M. M., Nair, P. H., ... Streicher, O. (2013). Astropy: A community Python package for astronomy. *Astronomy and Astrophysics*, 558, A33. <https://doi.org/10.1051/0004-6361/201322068>
- Barbary, K. (2016a). *Extinction v0.3.0*. Zenodo. <https://doi.org/10.5281/zenodo.804967>
- Barbary, K. (2016b). SEP: Source extractor as a library. *Journal of Open Source Software*, 1(6), 58. <https://doi.org/10.21105/joss.00058>
- Betoule, M., Kessler, R., Guy, J., Mosser, J., Hardin, D., Biswas, R., Astier, P., El-Hage, P., Konig, M., Kuhlmann, S., Mennier, J., Pain, R., Regnault, N., Balland, C., Bassett, B. A., Brown, P. J., Campbell, H., Carlberg, R. G., Cellier-Holzem, F., ... Wheeler, C. J. (2014). Improved cosmological constraints from a joint analysis of the SDSS-II and SNLS supernova samples. *Astronomy and Astrophysics*, 568, A22. <https://doi.org/10.1051/0004-6361/201423413>
- Bradley, L., Sipőcz, B., Robitaille, T., Tollerud, E., Vinícius, Z., Deil, C., Barbary, K., Wilson, T. J., Busko, I., Donath, A., Günther, H. M., Cara, M., Krachyon, Conseil, S., Bostroem, A., Droettboom, M., Bray, E. M., Lim, P. L., Bratholm, L. A., ... Souchereau, H. (2021). *Astropy/photutils: 1.3.0* (Version 1.3.0) [Computer software]. Zenodo. <https://doi.org/10.5281/zenodo.5796924>
- Brout, D., Sako, M., Scolnic, D., Kessler, R., D'Andrea, C. B., Davis, T. M., Hinton, S. R., Kim, A. G., Lasker, J., Macaulay, E., Möller, A., Nichol, R. C., Smith, M., Sullivan, M., Wolf, R. C., Allam, S., Bassett, B. A., Brown, P., Castander, F. J., ... DES COLLABORATION. (2019). First Cosmology Results Using Type Ia Supernovae from the Dark Energy Survey: Photometric Pipeline and Light-curve Data Release. *The Astrophysical Journal*, 874(1), 106. <https://doi.org/10.3847/1538-4357/ab06c1>
- Gaia Collaboration. (2016). The Gaia mission. *Astronomy and Astrophysics*, 595. <https://doi.org/10.1051/0004-6361/201629272>
- Ginsburg, A., Sipőcz, B. M., Brasseur, C. E., Cowperthwaite, P. S., Craig, M. W., Deil, C., Guillochon, J., Guzman, G., Liedtke, S., Lian Lim, P., Lockhart, K. E., Mommert, M., Morris, B. M., Norman, H., Parikh, M., Persson, M. V., Robitaille, T. P., Segovia, J.-C., Singer, L. P., ... a subset of the astropy collaboration. (2019). astroquery: An Astronomical Web-querying Package in Python. *The Astronomical Journal*, 157, 98. <https://doi.org/10.3847/1538-3881/aafc33>
- Graham, M., Plante, R., Tody, D., & Fitzpatrick, M. (2014). *PyVO: Python access to the Virtual Observatory* (p. ascl:1402.004). Astrophysics Source Code Library, record ascl:1402.004.
- Harris, C. R., Millman, K. J., Walt, S. J. van der, Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., Kerkwijk, M. H. van, Brett, M., Haldane, A., Río, J. F. del, Wiebe, M., Peterson, P., ... Oliphant, T. E. (2020). Array programming with NumPy. *Nature*, 585(7825), 357–362. <https://doi.org/10.1038/s41586-020-2649-2>
- Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. *Computing in Science Engineering*, 9(3), 90–95.
- Kelly, P. L., & others. (2010). Hubble Residuals of Nearby Type Ia Supernovae are Correlated with Host Galaxy Masses. *The Astrophysical Journal*, 715, 743–756. <https://doi.org/10.1088/0004-637X/715/2/743>
- Kelsey, L., Sullivan, M., Smith, M., Wiseman, P., Brout, D., Davis, T. M., Frohmaier, C., Galbany, L., Grayling, M., Gutiérrez, C. P., Hinton, S. R., Kessler, R., Lidman, C., Möller, A., Sako, M., Scolnic, D., Uddin, S. A., Vincenzi, M., Abbott, T. M. C., ... DES Collaboration. (2021). The effect of environment on Type Ia supernovae in the Dark Energy Survey

- three-year cosmological sample. *Monthly Notices of the Royal Astronomical Society*, 501(4), 4861–4876. <https://doi.org/10.1093/mnras/staa3924>
- Lampeitl, H., Nichol, R. C., Seo, H.-J., Giannantonio, T., Shapiro, C., Bassett, B., Percival, W. J., Davis, T. M., Dilday, B., Frieman, J., Garnavich, P., Sako, M., Smith, M., Sollerman, J., Becker, A. C., Cinabro, D., Filippenko, A. V., Foley, R. J., Hogan, C. J., ... Zheng, C. (2010). First-year Sloan Digital Sky Survey-II supernova results: consistency and constraints with other intermediate-redshift data sets. *Monthly Notices of the Royal Astronomical Society*, 401(4), 2331–2342. <https://doi.org/10.1111/j.1365-2966.2009.15851.x>
- McKinney, W., & others. (2010). Data structures for statistical computing in python. *Proceedings of the 9th Python in Science Conference*, 445, 51–56.
- Perlmutter, S., Aldering, G., Goldhaber, G., Knop, R. A., Nugent, P., Castro, P. G., Deustua, S., Fabbro, S., Goobar, A., Groom, D. E., Hook, I. M., Kim, A. G., Kim, M. Y., Lee, J. C., Nunes, N. J., Pain, R., Pennypacker, C. R., Quimby, R., Lidman, C., ... Project, T. S. C. (1999). Measurements of  $\Omega$  and  $\Lambda$  from 42 High-Redshift Supernovae. *The Astrophysical Journal*, 517(2), 565–586. <https://doi.org/10.1086/307221>
- Phillips, M. M. (1993). The Absolute Magnitudes of Type IA Supernovae. *Astrophysical Journal Letters*, 413, L105. <https://doi.org/10.1086/186970>
- Riess, A. G., Filippenko, A. V., Challis, P., Clocchiatti, A., Diercks, A., Garnavich, P. M., Gilliland, R. L., Hogan, C. J., Jha, S., Kirshner, R. P., Leibundgut, B., Phillips, M. M., Reiss, D., Schmidt, B. P., Schommer, R. A., Smith, R. C., Spyromilio, J., Stubbs, C., Suntzeff, N. B., & Tonry, J. (1998). Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant. *The Astronomical Journal*, 116(3), 1009–1038. <https://doi.org/10.1086/300499>
- Rigault, M., Copin, Y., Aldering, G., Antilogus, P., Aragon, C., Bailey, S., Baltay, C., Bongard, S., Buton, C., Canto, A., Cellier-Holzem, F., Childress, M., Chotard, N., Fakhouri, H. K., Feindt, U., Fleury, M., Gangler, E., Greskovic, P., Guy, J., ... Weaver, B. A. (2013). Evidence of environmental dependencies of Type Ia supernovae from the Nearby Supernova Factory indicated by local  $H\alpha$ . *Astronomy and Astrophysics*, 560, A66. <https://doi.org/10.1051/0004-6361/201322104>
- Roman, M., Hardin, D., Betoule, M., Astier, P., Ballew, C., Ellis, R. S., Fabbro, S., Guy, J., Hook, I., Howell, D. A., Lidman, C., Mitra, A., Möller, A., Mourão, A. M., Neveu, J., Palanque-Delabrouille, N., Pritchard, C. J., Regnault, N., Ruhlmann-Kleider, V., ... Sullivan, M. (2018). Dependence of Type Ia supernova luminosities on their local environment. *Astronomy and Astrophysics*, 615, A68. <https://doi.org/10.1051/0004-6361/201731425>
- Scolnic, D. M., Jones, D. O., Rest, A., Pan, Y. C., Chornock, R., Foley, R. J., Huber, M. E., Kessler, R., Narayan, G., Riess, A. G., Rodney, S., Berger, E., Brout, D. J., Challis, P. J., Drout, M., Finkbeiner, D., Lunnan, R., Kirshner, R. P., Sanders, N. E., ... Smith, K. W. (2018). The Complete Light-curve Sample of Spectroscopically Confirmed SNe Ia from Pan-STARRS1 and Cosmological Constraints from the Combined Pantheon Sample. *The Astrophysical Journal*, 859(2), 101. <https://doi.org/10.3847/1538-4357/aab9bb>
- Sullivan, M., Conley, A., Howell, D. A., Neill, J. D., Astier, P., Ballew, C., Basa, S., Carlberg, R. G., Fouchez, D., Guy, J., Hardin, D., Hook, I. M., Pain, R., Palanque-Delabrouille, N., Perrett, K. M., Pritchard, C. J., Regnault, N., Rich, J., Ruhlmann-Kleider, V., ... Walker, E. S. (2010). The dependence of Type Ia Supernovae luminosities on their host galaxies. *Monthly Notices of the Royal Astronomical Society*, 406(2), 782–802. <https://doi.org/10.1111/j.1365-2966.2010.16731.x>
- Tripp, R. (1998). A two-parameter luminosity correction for Type IA supernovae. *Astronomy and Astrophysics*, 331, 815–820.