

QSONic: fast quasar continuum fitting

Naim Göksel Karaçaylı  1,2,3

1 Center for Cosmology and AstroParticle Physics, The Ohio State University, 191 West Woodruff Avenue, Columbus, OH 43210, USA 2 Department of Astronomy, The Ohio State University, 4055 McPherson Laboratory, 140 W 18th Avenue, Columbus, OH 43210, USA 3 Department of Physics, The Ohio State University, 191 West Woodruff Avenue, Columbus, OH 43210, USA

DOI: [10.21105/joss.06373](https://doi.org/10.21105/joss.06373)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: Ivelina Momcheva  

Reviewers:

- [@gcalderone](#)
- [@corentinravoux](#)

Submitted: 18 January 2024

Published: 08 May 2024

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

Distant quasars show absorption features at wavelengths shorter than the Ly α emission line in their spectrum due to intervening neutral hydrogen gas. The Ly α forest technique uses these absorption lines to map out the matter distribution in vast volumes when the universe was 1.5–3 billion years old ($2 \lesssim z \lesssim 5$). This map of the cosmos has successfully constrained cosmological models using the Baryon Acoustic Oscillations feature in the 3D correlation function ([du Mas des Bourboux et al., 2020](#)), various dark matter models ([Armengaud et al., 2017; Iršič et al., 2017](#)) and the sum of the neutrino masses ([Palanque-Delabrouille et al., 2015](#)) using the 1D Ly α forest power spectrum ([Karaçaylı et al., 2022, 2024; Ravoux et al., 2023](#)).

The density of the intervening gas is inferred from the transmitted flux fraction, which is the absorption depth relative to the unabsorbed quasar continuum. Recent studies have devised a simple continuum model to homogenize the analysis across quasars so that the errors in the estimated continuum are contained to well-known modes, such as the amplitude and slope of the quasar continuum. This model assumes each quasar continuum C_q is a scaling of a mean quasar continuum shape \bar{C} by a polynomial of $\ln \lambda$. For example, an order 1 polynomial introduces amplitude a_q and slope b_q terms for each quasar such that $C_q = (a_q + b_q \ln \lambda) \bar{C}$. The quasar continuum fitting algorithm is then an iterative solution that minimizes χ^2 for each quasar and updates the mean quasar continuum using the stack of all quasar continua after every iteration ([du Mas des Bourboux et al., 2020; Karaçaylı et al., 2024](#)). This method requires efficient parallelization due to the computational time and memory needed.

Statement of need

QSONic is an MPI-parallelized, highly optimized Python package specifically built for the Dark Energy Spectroscopic Instrument (DESI), which will collect approximately one million Ly α quasar spectra in the near future ([DESI Collaboration et al., 2016](#)). QSONic is designed to overcome the numerical challenge posed by this large quantity of spectra by MPI parallelization that can distribute the memory and workload across multiple compute nodes. Each parallel process is responsible for independent sky regions based on HEALPix ([Górski et al., 2005](#)). Reading the data and fitting the continuum are done in parallel. After all quasars are fit with a continuum, all processes synchronize to calculate the mean continuum and check for convergence.

Furthermore, QSONic provides an efficient API to read and manipulate DESI quasar spectra to allow implementation of other algorithms within the Ly α forest framework. It saves detailed intermediate data products including best-fit parameters for each quasar and the covariance matrix used in the pipeline noise calibration correction (η parameter in references, [du Mas des](#)

Bourboux et al., 2020; Karaçaylı et al., 2024). These intermediate products are crucial to ascertain the precision of the fitted continua and the pipeline noise corrections.

QS0nic is built on the same principle of `picca` continuum fitting algorithm. Unlike `picca`, QS0nic has robust convergence criteria, provides a generic API for DESI quasar spectra, and saves more intermediate data products. Due to optimized IO routines, MPI parallelization and efficient numerical algorithms, it may be faster than `picca`. DESI Ly α forest working group and its ancillary topical groups are responsible for coordinating the efforts between different packages.

QS0nic will be used in future scientific publications of the 1D Ly α forest power spectrum and the 3D correlation function measurements from DESI data.

References

- Armengaud, E., Palanque-Delabrouille, N., Yèche, C., Marsh, D. J. E., & Baur, J. (2017). Constraining the mass of light bosonic dark matter using SDSS Lyman- α forest. *Monthly Notices of the Royal Astronomical Society*, 471(4), 4606–4614. <https://doi.org/10.1093/mnras/stx1870>
- DESI Collaboration, Aghamousa, A., Aguilar, J., Ahlen, S., Alam, S., Allen, L. E., Allende Prieto, C., Annis, J., Bailey, S., Balland, C., Ballester, O., Baltay, C., Beaufore, L., Bebek, C., Beers, T. C., Bell, E. F., Bernal, J. L., Besuner, R., Beutler, F., ... Zu, Y. (2016). The DESI Experiment Part I: Science, Targeting, and Survey Design. *arXiv e-Prints*, arXiv:1611.00036. <https://doi.org/10.48550/arXiv.1611.00036>
- du Mas des Bourboux, H., Rich, J., Font-Ribera, A., de Sainte Agathe, V., Farr, J., Etourneau, T., Le Goff, J.-M., Cuceu, A., Balland, C., Bautista, J. E., Blomqvist, M., Brinkmann, J., Brownstein, J. R., Chabanier, S., Chaussidon, E., Dawson, K., González-Morales, A. X., Guy, J., Lyke, B. W., ... Youles, S. (2020). The Completed SDSS-IV Extended Baryon Oscillation Spectroscopic Survey: Baryon Acoustic Oscillations with Ly α Forests. *Astrophysical Journal*, 901(2), 153. <https://doi.org/10.3847/1538-4357/abb085>
- Górski, K. M., Hivon, E., Banday, A. J., Wandelt, B. D., Hansen, F. K., Reinecke, M., & Bartelmann, M. (2005). HEALPix: A Framework for High-Resolution Discretization and Fast Analysis of Data Distributed on the Sphere. *Astrophysical Journal*, 622(2), 759–771. <https://doi.org/10.1086/427976>
- Iršič, V., Viel, M., Haehnelt, M. G., Bolton, J. S., Cristiani, S., Becker, G. D., D'Odorico, V., Cupani, G., Kim, T.-S., Berg, T. A. M., López, S., Ellison, S., Christensen, L., Denney, K. D., & Worseck, G. (2017). New constraints on the free-streaming of warm dark matter from intermediate and small scale Lyman- α forest data. *Phys. Rev. D.*, 96(2), 023522. <https://doi.org/10.1103/PhysRevD.96.023522>
- Karaçaylı, N. G., Martini, P., Guy, J., Ravoux, C., Karim, M. L. A., Armengaud, E., Walther, M., Aguilar, J., Ahlen, S., Bailey, S., Bautista, J., Beltran, S. F., Brooks, D., Cabayol-Garcia, L., Chabanier, S., Chaussidon, E., Chaves-Montero, J., Dawson, K., Cruz, R. de la, ... Zhou, Z. (2024). Optimal 1D Ly α Forest Power Spectrum Estimation – III. DESI early data. *Monthly Notices of the Royal Astronomical Society*, stae171. <https://doi.org/10.1093/mnras/stae171>
- Karaçaylı, N. G., Padmanabhan, N., Font-Ribera, A., Iršič, V., Walther, M., Brooks, D., Gaztañaga, E., Kehoe, R., Levi, M., Ntelis, P., Palanque-Delabrouille, N., & Tarlé, G. (2022). Optimal 1D Ly α forest power spectrum estimation - II. KODIAQ, SQUAD, and XQ-100. *Monthly Notices of the Royal Astronomical Society*, 509(2), 2842–2855. <https://doi.org/10.1093/mnras/stab3201>
- Palanque-Delabrouille, N., Yèche, C., Lesgourgues, J., Rossi, G., Borde, A., Viel, M., Aubourg, E., Kirkby, D., LeGoff, J.-M., Rich, J., Roe, N., Ross, N. P., Schneider, D. P., &

- Weinberg, D. (2015). Constraint on neutrino masses from SDSS-III/BOSS Ly α forest and other cosmological probes. *J. Cosmology Astropart. Phys.*, 2015(2), 045–045. <https://doi.org/10.1088/1475-7516/2015/02/045>
- Ravoux, C., Abdul Karim, M. L., Armengaud, E., Walther, M., Karaçaylı, N. G., Martini, P., Guy, J., Aguilar, J. N., Ahlen, S., Bailey, S., Bautista, J., Beltran, S. F., Brooks, D., Cabayol-Garcia, L., Chabanier, S., Chaussidon, E., Chaves-Montero, J., Dawson, K., de la Cruz, R., ... Zhou, Z. (2023). The Dark Energy Spectroscopic Instrument: one-dimensional power spectrum from first Ly α forest samples with Fast Fourier Transform. *Monthly Notices of the Royal Astronomical Society*, 526(4), 5118–5140. <https://doi.org/10.1093/mnras/stad3008>