






# TopologicalNumbers.jl: A Julia package for topological number computation

Keisuke Adachi <sup>1,2\*</sup> and Minoru Kanega <sup>2\*</sup>

<sup>1</sup> Department of Physics, Ibaraki University, Mito, Ibaraki, Japan <sup>2</sup> Department of Physics, Chiba University, Chiba, Japan ¶ Corresponding author \* These authors contributed equally.

DOI: [10.21105/joss.06944](https://doi.org/10.21105/joss.06944)

## Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Olexandr Kononov](#) 

## Reviewers:

- [@lf28](#)
- [@michiexile](#)

Submitted: 26 January 2024

Published: 23 April 2025

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

## Summary

TopologicalNumbers.jl is an open-source Julia package designed to calculate topological invariants — mathematical quantities that characterize the properties of materials in condensed matter physics. These invariants, such as the Chern number and the  $\mathbb{Z}_2$  invariant, are crucial for understanding exotic materials like topological insulators and superconductors, which have potential applications in advanced electronics, spintronics, and quantum computing ([Hasan & Kane, 2010](#); [Nayak et al., 2008](#); [Qi & Zhang, 2011](#)). This package provides researchers and educators with an easy-to-use and efficient toolset to compute these invariants across various dimensions and symmetry classes, facilitating the exploration and discovery of new topological phases of matter.

## Statement of need

Understanding the properties of materials is essential in solid-state physics. For example, electrical conductivity is a key physical quantity indicating how well a material conducts electric current. Typically, when a weak electric field is applied to a material, if quantum eigenstates exist in the bulk into which electrons can transition, the material exhibits finite electrical conductivity and behaves as a metal. Conversely, if such states do not exist, the electrical conductivity is low, and the material behaves as an insulator. Since the 1980s, it has been revealed that certain materials exhibit states in which the bulk is insulating while the material's surface has conducting electronic states ([Hasan & Kane, 2010](#); [Qi & Zhang, 2011](#)). These materials are known as topological electronic systems, including quantum Hall insulators and topological insulators. Due to these novel properties, extensive research has been conducted to identify candidate materials and evaluate their characteristics.

The features of surface conducting states are determined by the topology of quantum eigenstates in momentum space. Topological numbers, such as the first Chern number, the second Chern number, and the  $\mathbb{Z}_2$  invariant, are used to characterize these properties ([Kane & Mele, 2005](#); [Thouless et al., 1982](#)). A typical example is the quantum Hall effect, where applying a weak electric field to a two-dimensional material results in a quantized finite electrical conductivity (Hall conductivity) perpendicular to the applied field ([Thouless et al., 1982](#)). The Hall conductivity  $\sigma_{xy}$  is characterized by the first Chern number  $\nu \in \mathbb{Z}$  and is given by  $\sigma_{xy} = \frac{e^2}{h} \nu$ , where  $e$  is the elementary charge and  $h$  is Planck's constant. Other topological numbers similarly serve as important physical quantities that characterize systems, depending on their dimensions and symmetry classes ([Ryu et al., 2010](#)).

Obtaining topological numbers often requires extensive numerical calculations, which may demand considerable computational effort before achieving convergence. Therefore, creating tools that simplify the computation of these quantities will advance research on topological

phases of matter. Several methods have been developed to enable efficient computation of certain topological numbers (Fukui et al., 2005; Fukui & Hatsugai, 2007; Mochol-Grzelak et al., 2018; Shiozaki, 2023). However, since each method is typically specialized for specific dimensions or symmetry classes, one must often implement algorithms separately for each problem. Our project, `TopologicalNumbers.jl`, aims to provide a package that can efficiently and easily compute topological numbers across various dimensions and symmetry classes.

This package currently includes several methods for calculating topological numbers. The first is the Fukui–Hatsugai–Suzuki (FHS) method for computing the first Chern number in two-dimensional solid-state systems (Fukui et al., 2005). The first Chern number is obtained by integrating the Berry curvature, derived from the Hamiltonian’s eigenstates, over the Brillouin zone. The FHS method enables efficient computation by discretizing the Berry curvature in the Brillouin zone. Several methods based on the FHS approach have been proposed to compute various topological numbers. One such method calculates the second Chern number in four-dimensional systems (Mochol-Grzelak et al., 2018). The  $\mathbb{Z}_2$  invariant can be computed in two-dimensional systems with time-reversal symmetry (Fukui & Hatsugai, 2007; Shiozaki, 2023). The FHS method also applies to identifying Weyl points and Weyl nodes in three-dimensional systems (Du et al., 2017; Hirayama et al., 2015, 2018; Yang et al., 2011).

Currently, there is no comprehensive Julia package that implements all these calculation methods. On other platforms, software packages using different approaches — such as those based on Wannier charge centers (Soluyanov, 2011) or Wilson loops (Yu et al., 2011) — are available. For example, `Z2Pack` (Gresch et al., 2017) is a Python-based tool widely used for calculating the  $\mathbb{Z}_2$  invariant and the first Chern number. `WannierTools` (Wu et al., 2018) offers powerful features for analyzing topological materials but is implemented in Fortran, which may pose a steep learning curve for some users.

`TopologicalNumbers.jl` distinguishes itself by providing an efficient, pure Julia implementation. Julia is known for its high performance and user-friendly syntax. This package supports various topological invariants across multiple dimensions and symmetry classes, including the first and second Chern numbers and the  $\mathbb{Z}_2$  invariant. It also offers parallel computing capabilities through `MPI.jl`, enhancing computational efficiency for large-scale problems. By leveraging Julia’s multiple dispatch feature, we adopt a consistent interface using the `Problem`, `Algorithm`, and `solve` style — similar to `DifferentialEquations.jl` (Rackauckas & Nie, 2017) — to improve extensibility. With these features, `TopologicalNumbers.jl` achieves a unique balance of performance, usability, maintainability, and extensibility, providing an alternative perspective rather than directly competing with other libraries.

Additionally, to compute the  $\mathbb{Z}_2$  invariant, which requires calculating the Pfaffian, we have ported `PFAPACK` to Julia. `PFAPACK` is a Fortran/C++/Python library for computing the Pfaffian of skew-symmetric matrices (Wimmer, 2012). Our package includes pure Julia implementations of all originally provided functions. While `SkewLinearAlgebra.jl` exists as an official Julia package for computing the Pfaffian of real skew-symmetric matrices, to our knowledge, `TopologicalNumbers.jl` is the first official package to offer a pure Julia implementation that handles complex skew-symmetric matrices.

## Usage

Users can easily compute topological numbers using the various methods included in this package. In the simplest case, they need only provide a function that returns the Hamiltonian matrix as a function of the wave numbers. Computations are performed by creating the corresponding `Problem` instance and calling the `solve` function (`solve(Problem)`). The package also provides the `calcPhaseDiagram` function, which enables the computation of topological numbers in one-dimensional or two-dimensional parameter spaces by specifying the `Problem` and parameter ranges (`calcPhaseDiagram(Problem, range...)`).

Moreover, utility functions such as `showBand`, `plot1D`, and `plot2D` are available for visualizing

energy band structures and phase diagrams. We also offer various model Hamiltonians, such as the Su–Schrieffer–Heeger (SSH) model (Su et al., 1979) and the Haldane model (Haldane, 1988), allowing users to quickly test functionalities and learn how to use these features.

## Acknowledgements

The authors are grateful to Takahiro Fukui for fruitful discussions. M. K. was supported by JST, the establishment of university fellowships towards the creation of science technology innovation (Grant No. JPMJFS2107), and by JST SPRING (Grant No. JPMJSP2109).

## References

- Du, Y., Bo, X., Wang, D., Kan, E., Duan, C.-G., Savrasov, S. Y., & Wan, X. (2017). Emergence of topological nodal lines and type-II Weyl nodes in the strong spin-orbit coupling system  $\text{InNbX}_2$  ( $x = \text{s, se}$ ). *Phys. Rev. B*, 96(23), 235152. <https://doi.org/10.1103/PhysRevB.96.235152>
- Fukui, T., & Hatsugai, Y. (2007). Quantum Spin Hall Effect in Three Dimensional Materials: Lattice Computation of  $\mathbb{Z}_2$  Topological Invariants and Its Application to Bi and Sb. *J. Phys. Soc. Jpn.*, 76(5), 053702. <https://doi.org/10.1143/JPSJ.76.053702>
- Fukui, T., Hatsugai, Y., & Suzuki, H. (2005). Chern Numbers in Discretized Brillouin Zone: Efficient Method of Computing (Spin) Hall Conductances. *J. Phys. Soc. Jpn.*, 74(6), 1674–1677. <https://doi.org/10.1143/JPSJ.74.1674>
- Gresch, D., Autès, G., Yazyev, O. V., Troyer, M., Vanderbilt, D., Bernevig, B. A., & Soluyanov, A. A. (2017). Z2Pack: Numerical implementation of hybrid Wannier centers for identifying topological materials. *Phys. Rev. B*, 95(7), 075146. <https://doi.org/10.1103/PhysRevB.95.075146>
- Haldane, F. D. M. (1988). Model for a Quantum Hall Effect without Landau Levels: Condensed-Matter Realization of the "Parity Anomaly". *Phys. Rev. Lett.*, 61(18), 2015–2018. <https://doi.org/10.1103/PhysRevLett.61.2015>
- Hasan, M. Z., & Kane, C. L. (2010). Colloquium: Topological insulators. *Rev. Mod. Phys.*, 82(4), 3045–3067. <https://doi.org/10.1103/RevModPhys.82.3045>
- Hirayama, M., Okugawa, R., Ishibashi, S., Murakami, S., & Miyake, T. (2015). Weyl Node and Spin Texture in Trigonal Tellurium and Selenium. *Phys. Rev. Lett.*, 114(20), 206401. <https://doi.org/10.1103/PhysRevLett.114.206401>
- Hirayama, M., Okugawa, R., & Murakami, S. (2018). Topological Semimetals Studied by Ab Initio Calculations. *J. Phys. Soc. Jpn.*, 87(4), 041002. <https://doi.org/10.7566/JPSJ.87.041002>
- Kane, C. L., & Mele, E. J. (2005).  $\mathbb{Z}_2$  Topological Order and the Quantum Spin Hall Effect. *Phys. Rev. Lett.*, 95(14), 146802. <https://doi.org/10.1103/PhysRevLett.95.146802>
- Mochol-Grzelak, M., Dauphin, A., Celi, A., & Lewenstein, M. (2018). Efficient algorithm to compute the second Chern number in four dimensional systems. *Quantum Sci. Technol.*, 4(1), 014009. <https://doi.org/10.1088/2058-9565/aae93b>
- Nayak, C., Simon, S. H., Stern, A., Freedman, M., & Das Sarma, S. (2008). Non-Abelian anyons and topological quantum computation. *Rev. Mod. Phys.*, 80(3), 1083–1159. <https://doi.org/10.1103/RevModPhys.80.1083>
- Qi, X.-L., & Zhang, S.-C. (2011). Topological insulators and superconductors. *Rev. Mod. Phys.*, 83(4), 1057–1110. <https://doi.org/10.1103/RevModPhys.83.1057>

- Rackauckas, C., & Nie, Q. (2017). DifferentialEquations.jl – A Performant and Feature-Rich Ecosystem for Solving Differential Equations in Julia. *J. Open Res. Softw.*, 5(1), 15–15. <https://doi.org/10.5334/jors.151>
- Ryu, S., Schnyder, A. P., Furusaki, A., & Ludwig, A. W. W. (2010). Topological insulators and superconductors: Tenfold way and dimensional hierarchy. *New J. Phys.*, 12(6), 065010. <https://doi.org/10.1088/1367-2630/12/6/065010>
- Shiozaki, K. (2023). A discrete formulation of the Kane-Mele  $\mathbb{Z}_2$  invariant (No. arXiv:2305.05615). arXiv. <https://doi.org/10.48550/arXiv.2305.05615>
- Soluyanov, A. A. (2011). Computing topological invariants without inversion symmetry. *Phys. Rev. B*, 83(23). <https://doi.org/10.1103/PhysRevB.83.235401>
- Su, W. P., Schrieffer, J. R., & Heeger, A. J. (1979). Solitons in Polyacetylene. *Phys. Rev. Lett.*, 42(25), 1698–1701. <https://doi.org/10.1103/PhysRevLett.42.1698>
- Thouless, D. J., Kohmoto, M., Nightingale, M. P., & den Nijs, M. (1982). Quantized Hall Conductance in a Two-Dimensional Periodic Potential. *Phys. Rev. Lett.*, 49(6), 405–408. <https://doi.org/10.1103/PhysRevLett.49.405>
- Wimmer, M. (2012). Algorithm 923: Efficient Numerical Computation of the Pfaffian for Dense and Banded Skew-Symmetric Matrices. *ACM Trans. Math. Softw.*, 38(4), 30:1–30:17. <https://doi.org/10.1145/2331130.2331138>
- Wu, Q., Zhang, S., Song, H.-F., Troyer, M., & Soluyanov, A. A. (2018). WannierTools: An open-source software package for novel topological materials. *Comput. Phys. Commun.*, 224, 405–416. <https://doi.org/10.1016/j.cpc.2017.09.033>
- Yang, K.-Y., Lu, Y.-M., & Ran, Y. (2011). Quantum Hall effects in a Weyl semimetal: Possible application in pyrochlore iridates. *Phys. Rev. B*, 84(7), 075129. <https://doi.org/10.1103/PhysRevB.84.075129>
- Yu, R., Qi, X. L., Bernevig, A., Fang, Z., & Dai, X. (2011). Equivalent expression of  $\mathbb{Z}_2$  topological invariant for band insulators using the non-Abelian Berry connection. *Phys. Rev. B*, 84(7), 075119. <https://doi.org/10.1103/PhysRevB.84.075119>