

GASTLI: A Python package for coupled interior–atmosphere modelling of volatile-rich planets

Lorena Acuña  ^{1¶}, Laura Kreidberg  ¹, Meng Zhai  ^{1,2}, Paul Mollière  ¹, and Morgan Fouesneau  ¹

¹ Max Planck Institute for Astronomy, Königstuhl 17, 69117 Heidelberg, Germany ² Chinese Academy of Sciences South America Center for Astronomy (CASSACA), National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100101, PR China ¶ Corresponding author

DOI: [10.21105/joss.07288](https://doi.org/10.21105/joss.07288)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Josh Borrow](#)  

Reviewers:

- [@ivalaginja](#)
- [@MartianColonist](#)

Submitted: 08 August 2024

Published: 21 March 2025

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

Gaseous exoplanets are far more diverse in mass, density and temperature than their Solar System counterparts. These range from temperate sub-Neptunes whose masses and sizes are below that of the Solar System ice giants, to extremely irradiated, Jupiter-mass exoplanets known as hot Jupiters. The bulk composition of sub-Neptunes and gas giants is key to understanding their formation and evolution pathways, including their dominant formation mechanism (Bitsch & Mah, 2023; Ginzburg & Chiang, 2020; Schneider & Bitsch, 2021; Teske et al., 2019; Thorngren et al., 2016) and interactions between the irradiation of the host star and the planetary atmosphere (Acuña et al., 2022; Owen & Wu, 2013; Rogers et al., 2024). Their building blocks consist of rocks and water (metals), and hydrogen and helium (H/He), which are present in a wide variety of mass fractions in the exoplanet population (Helled et al., 2022). For exoplanets that contain H/He, the planetary radius (or density) not only depends on mass, but also on the temperature due to the distance from its host star (Burrows et al., 1997), its age (Baraffe et al., 2003), the mass of the core (Baraffe et al., 2008), and the fraction of H/He in its envelope (Rogers et al., 2023). The inference of bulk composition from observational data – mass, radius, equilibrium temperature, age and atmospheric metallicity – requires non-trivial numerical methods that solve the equations of planetary interior structure (Miguel & Vazan, 2023), in addition to Bayesian methods to solve the inverse problem (Acuña et al., 2021; Bloot et al., 2023; Dorn et al., 2015; Otegi et al., 2020). Additionally, interior models need to incorporate the effects of processes occurring in the planet atmosphere, such as radiative transfer (Baraffe et al., 2003; Marley et al., 2007), and the thermodynamical properties of the different materials, provided by the equation of state (EOS, Acuña et al., 2024; Howard et al., 2023; Howard & Guillot, 2023).

GASTLI (GAS gianT modeL for Interiors) is a Python package for interior structure modelling of volatile-rich exoplanets. The package combines a Fortran back end with a Python-based user interface. This enables fast convergence of the interior structure to a solution that fulfills the user input, while having a flexible and user-friendly interface. GASTLI was specifically designed to compute a wider range of combinations of core mass and H/He content in the envelope compared to existing open-source interior models. GASTLI also incorporates state-of-the-art thermodynamical data and EOSs for rock, water and H/He. One of its key features is the coupling class, which couples the interior structure model to a grid of atmospheric temperature profiles to determine self-consistently the temperature of the outermost layer. GASTLI is released with a default atmospheric grid that assumes equilibrium chemistry and a clear atmosphere at varying internal temperatures computed with petitCODE (Mollière et al., 2015, 2017). Nonetheless, the flexible interface enables the user to use their own custom atmospheric grid, which is useful to explore the effect of the processes assumed in the upper atmosphere in the interior, such as the type of chemistry (equilibrium or disequilibrium), cloud

implementations or 3D global circulation (GCM).

Statement of need

Interior structure codes are essential to carry out planet composition analyses and Bayesian inverse fitting—known as retrievals—of mass and radius data. MESA (Paxton et al., 2011, 2013, 2015, 2018, 2019) is a well-known stellar interior structure software that has been applied to sub-Neptunes and gas giants (Chen & Rogers, 2016; Müller & Helled, 2021). The compositions of the envelope in MESA are limited to low metal contents due to the default EOSs, which is particularly limiting for planets with Neptune's mass or lower. The default MESA EOS can incorporate up to a 4x solar composition in the envelope. For reference, Saturn's upper atmosphere is approximately 10x solar (Atreya et al., 2022), while Neptune's measured envelope metallicity is 80x solar (Irwin et al., 2021; Karkoschka & Tomasko, 2011). Therefore, the implementation of alternative EOS in MESA are necessary, which is not straightforward for the user given the complexity of the code, and the input system based on Fortran namelist files – known as inlists. These input namelists allow users to specify initial conditions, control settings, and adjust physical parameters within the MESA software. MESA requires at least one minute of computational time for a single forward model calculation, making it difficult to couple with atmospheric models. This coupling is important because atmospheric conditions can significantly influence the the output of the interior calculations (Linder et al., 2019; Lopez & Fortney, 2014; Poser et al., 2019; Poser & Redmer, 2024). MAGRATHEA (Huang et al., 2022) is another open-source planet interior model that is a computationally fast (< 2 seconds for a single forward model) and based on C++. Its layer structure (ideal gas EOS on top of liquid/ice water layer) makes its adaptation for applications to sub-Neptunes and gas giants a challenge to new users because these types of planet require the treatment of nonideal, high-pressure H/He (Chabrier & Debras, 2021; Saumon et al., 1995) and metal (water, Haldemann et al., 2020; Mazevet et al., 2019; Thomas & Madhusudhan, 2016) envelopes. Additionally, Exolnt (Wang et al., 2019) and ExoPlex (Unterborn et al., 2023) are interior structure model packages that include Fe-rich cores and rock mantles dedicated to planets without volatile gas species, such as super-Earths and rocky planets ($R < 2 R_{\oplus}$).

The GASTLI Python package

The interior model module is contained in the `GASTLI.int_planet` class. The interior structure equations for hydrostatic equilibrium, Gauss's theorem for gravitational acceleration, convection, and mass conservation are solved by combining integration by the trapezoidal rule with an iteration scheme over all profiles (pressure, temperature, gravity and density). In each iteration, the trapezoidal integrations are carried for each point in a 1D grid that represents the radius, from the center of the planet to the outermost boundary of the top layer. The density is calculated by evaluating the EOS for rock (Lyon, 1992; Miguel et al., 2022), water (Mazevet et al., 2019) and H/He (Chabrier & Debras, 2021; Howard & Guillot, 2023) at the pressure and temperature conditions of each point in the spatial grid. The iteration sequence is considered to have converged when the difference between iterations is less than a given precision that can be modified by the user. The recommended and default value for this precision is 10^{-5} . The iteration scheme can also be stopped at a given number of iterations defined by the user, with a default of 30 maximum iterations, and a maximum possible value of 99 iterations. The boundary conditions to solve these equations include the top layer's pressure and temperature, which can be defined by the user. In addition to the profile convergence, the interior module ensures that the surface boundary conditions and the input mass are satisfied. A complete description of the interior module integration scheme can be found in Brugger (2018), and has been widely used (Acuña et al., 2021, 2022, 2023; Aguichine et al., 2021; Brugger et al., 2016, 2017; Mousis et al., 2020; Vivien et al., 2022). The new features of the interior module include the optimization of several of its routines, such as the trapezoidal integration and the

evaluation of the EOS's density tables, and the parallelization of the calculation of the interior structure profiles with OpenMP in Fortran. One single computation of the interior module (function `calc_radius` in the `GASTLI.int_planet` class) for the default precision and number of iterations takes 1.89 seconds in an Apple M2 CPU, with parallelization on 8 cores (four cores operating at 3.49 GHz and the other four cores operating at 2.42 GHz).

The class `atm_models_interp` encloses the function that interpolates the atmospheric grid to obtain the temperature profile and interior boundary temperature. In addition, this class also provides the functions that calculate the atmospheric thickness and the atmosphere density profile. The latter is obtained by evaluating the EOS used by the interior module for H/He, and the AQUA water EOS (Haldemann et al., 2020). The atmospheric grid class and the interior module class are then combined in the Coupling class. This class self-consistently couples the interior module and the atmosphere grids to calculate the boundary temperature of the outermost layer of the interior model. See Acuña et al. (2021) for a detailed description of the coupling algorithm, which has been used in previous work (Acuña et al., 2022, 2023; Aguichine et al., 2021). Finally, the class `Thermal_evolution` contains the functions necessary to evaluate a sequence of coupled interior-atmosphere models at different internal temperatures and solve the luminosity equation to obtain the internal temperatures and radii as a function of age (Fortney et al., 2013; Poser et al., 2019). The complete validation of GASTLI against mass-radius relationships obtained with interior models used in previous work (Fortney et al., 2013; Müller & Helled, 2021) can be found in Acuña et al. (2024).




Figure 1: Schematic diagram of GASTLI's compositional structure, inputs and outputs.

In an interior retrieval, we fit an interior composition model to data, particularly to mass, radius, age, atmospheric metallicity and/or internal temperature. This is usually done through Bayesian model fitting, such as Markov chain Monte Carlo (Mosegaard & Tarantola, 1995) or nested sampling (Buchner, 2023). The likelihood function in interior retrievals is computed as a function of the squared residuals between the compositional model and the observed values of mass, radius, etc. The likelihood function used in interior retrievals can be found in equations 6 and 14 of Dorn et al. (2015) and Acuña et al. (2021), respectively. GASTLI can generate a grid of forward models that contains thousands of thermal evolution tracks and explores all possible parameter space for an exoplanet. For example, a grid of 4000 models can be computed in 12 hours, depending on the computer architecture and number of threads/cores available for parallelization, given that each model's computational time is 2-10 seconds. In GASTLI's documentation, we show an example of how to generate a grid and interpolate it

to carry an interior retrieval. This example uses emcee as a sampler ([Foreman-Mackey et al., 2013](#)).

Finally, GASTLI provides a default atmospheric grid for gas giants and sub-Neptunes with equilibrium temperatures under 1000 K, and atmospheric compositions from sub-solar to 250x solar (80% metal mass fraction). Our default grid assumes clear atmospheres in chemical equilibrium. If this grid is not appropriate for a particular target, the user may choose from a variety of model grids that are publicly available. These grids include clear, self-luminous atmospheres in equilibrium ([Linder et al., 2019](#); [Marley et al., 2021](#); [Samland et al., 2017](#)), clear, self-luminous in disequilibrium ([Karalidi et al., 2021](#); [Mukherjee et al., 2024](#)), cloudy self-luminous ([Jørgensen et al., 2024](#); [Lacy & Burrows, 2023](#); [Morley et al., 2024](#)), cloudy irradiated gas giants ([Charnay et al., 2018](#)), hot Jupiters ([Mollière et al., 2015](#); [Roth et al., 2024](#)) and temperate sub-Neptunes ([Kempton et al., 2023](#)). GASTLI's documentation includes an example of the grid file format and input.

Similar tools

The following table lists public software tools that provide mass-radius curves and possibly thermal evolution tracks. We distinguish between open-source interior structure packages, and neural networks together with grid interpolators. The latter may use tables and data generated with proprietary interior structure model codes.

Name and link	Type of software	Keywords	Reference
MESA	Interior structure model	Stars, hot Jupiters, gas giants; H/He and metals	Paxton et al. (2011), Paxton et al. (2013), Paxton et al. (2015), Paxton et al. (2018), Paxton et al. (2019)
Magrathea	Interior structure model	Super-Earths, rocky planets; liquid water	Huang et al. (2022)
ExoInt	Interior structure model	Dry super-Earths, rocky planets; Fe, rock	Wang et al. (2019)
ExoPlex	Interior structure model	Dry super-Earths, rocky planets; Fe, rock	Unterborn et al. (2023)
plaNETic	Trained neural network	Dry super-Earths, rocky planets; H/He, water, rock, Fe	Egger et al. (2024), Haldemann et al. (2024)
ExoMDN	Trained neural network	Super-Earths, sub-Neptunes; Fe, rock, liquid water, H/He. Retrievals only.	Baumeister & Tosi (2023)
Exoplanet Composition Interpolator	Grid interpolation (web interface)	Sub-Neptunes; H/He, rock, Fe	Lopez & Fortney (2014)

Name and link	Type of software	Keywords	Reference
HARDCORE	Grid interpolation (web interface)	Dry super-Earths, rocky planets; Fe, rock	Suissa et al. (2018)
planetsynth	Grid interpolator	Gas giants; H/He, metals	Müller & Helled (2021)
SMINT	Grid interpolator	Sub-Neptunes; H/He, water, rock, Fe	Piaulet et al. (2021); Tables from Lopez & Fortney (2014), Zeng et al. (2016), Zeng et al. (2019), Aguichine et al. (2021)
mr-plotter	Grid interpolator	Super-Earths, sub-Neptunes; H/He, water, rock, Fe	Castro-González et al. (2023); Tables from Lopez & Fortney (2014), Zeng et al. (2016), Zeng et al. (2019), Aguichine et al. (2021), Seager et al. (2007), Turbet et al. (2020), Haldemann et al. (2024)
Mardigras	Grid interpolator	Sub-Neptunes; Water, rock, Fe	Aguichine (2024); Tables from Lopez & Fortney (2014), Zeng et al. (2019), Aguichine et al. (2021)

Documentation

Users can find GASTLI's documentation [here](#).

Acknowledgements

We thank Yamila Miguel for sharing the reformatted EOS tables for silicate (dry sand) from the SESAME database. We acknowledge the support of the Data Science Group at the Max Planck Institute for Astronomy (MPIA) and especially Ivelina Momcheva for her invaluable assistance in optimizing the software and developing the GASTLI Python package. We thank Magali Deleuil, Olivier Mousis, Artyom Aguichine and Bastien Brugger for their contributions to the development of Marseille's Super-Earth Interior (MSEI) model, on which GASTLI's development is based. We thank the two JOSS reviewers, Iva Laginja and Ryan MacDonald, for their suggestions to improve the manuscript, as well as the JOSS editor, Josh Borrow.

References

- Acuña, L., Deleuil, M., & Mousis, O. (2023). Interior-atmosphere modelling to assess the observability of rocky planets with JWST. *Astronomy & Astrophysics*, 677, A14. <https://doi.org/10.1051/0004-6361/202245736>
- Acuña, L., Deleuil, M., Mousis, O., Marcq, E., Levesque, M., & Aguichine, A. (2021). Characterisation of the hydrospheres of TRAPPIST-1 planets. *Astronomy & Astrophysics*, 647, A53. <https://doi.org/10.1051/0004-6361/202039885>
- Acuña, L., Kreidberg, L., Zhai, M., & Mollière, P. (2024). GASTLI. An open-source coupled interior-atmosphere model to unveil gas-giant composition. *Astronomy & Astrophysics*, 688, A60. <https://doi.org/10.1051/0004-6361/202450559>

- Acuña, L., Lopez, T. A., Morel, T., Deleuil, M., Mousis, O., Aguichine, A., Marcq, E., & Santerne, A. (2022). Water content trends in K2-138 and other low-mass multi-planetary systems. *Astronomy & Astrophysics*, 660, A102. <https://doi.org/10.1051/0004-6361/202142374>
- Aguichine, A. (2024). mardigras: A visualization tool of theoretical mass–radius relations in the context of planetary science. *Research Notes of the American Astronomical Society*, 8(8), 216. <https://doi.org/10.3847/2515-5172/ad7506>
- Aguichine, A., Mousis, O., Deleuil, M., & Marcq, E. (2021). Mass-radius relationships for irradiated ocean planets. *The Astrophysical Journal*, 914(2), 84. <https://doi.org/10.3847/1538-4357/abfa99>
- Atreya, S. K., Crida, A., Guillot, T., Li, C., Lunine, J. I., Madhusudhan, N., Mousis, O., & Wong, M. H. (2022). The origin and evolution of Saturn: A post-Cassini perspective. *arXiv e-Prints*, arXiv:2205.06914. <https://doi.org/10.48550/arXiv.2205.06914>
- Baraffe, I., Chabrier, G., & Barman, T. (2008). Structure and evolution of super-Earth to super-Jupiter exoplanets. I. Heavy element enrichment in the interior. *Astronomy & Astrophysics*, 482(1), 315–332. <https://doi.org/10.1051/0004-6361:20079321>
- Baraffe, I., Chabrier, G., Barman, T. S., Allard, F., & Hauschildt, P. H. (2003). Evolutionary models for cool brown dwarfs and extrasolar giant planets. The case of HD 209458. *Astronomy & Astrophysics*, 402, 701–712. <https://doi.org/10.1051/0004-6361:20030252>
- Baumeister, P., & Tosi, N. (2023). ExoMDN: Rapid characterization of exoplanet interior structures with mixture density networks. *Astronomy & Astrophysics*, 676, A106. <https://doi.org/10.1051/0004-6361/202346216>
- Bitsch, B., & Mah, J. (2023). Enriching inner discs and giant planets with heavy elements. *Astronomy & Astrophysics*, 679, A11. <https://doi.org/10.1051/0004-6361/202347419>
- Bloot, S., Miguel, Y., Bazot, M., & Howard, S. (2023). Exoplanet interior retrievals: Core masses and metallicities from atmospheric abundances. *Monthly Notices of the Royal Astronomical Society*, 523(4), 6282–6292. <https://doi.org/10.1093/mnras/stad1873>
- Brugger, B. (2018). *Structure interne et minéralogie des exoplanètes terrestres de faible masse* [PhD thesis]. <http://www.theses.fr/2018AIXM0324/document>
- Brugger, B., Mousis, O., Deleuil, M., & Deschamps, F. (2017). Constraints on super-Earth interiors from stellar abundances. *The Astrophysical Journal*, 850(1), 93. <https://doi.org/10.3847/1538-4357/aa965a>
- Brugger, B., Mousis, O., Deleuil, M., & Lunine, J. I. (2016). Possible internal structures and compositions of Proxima Centauri b. *The Astrophysical Journal Letters*, 831(2), L16. <https://doi.org/10.3847/2041-8205/831/2/L16>
- Buchner, J. (2023). Nested sampling methods. *Statistics Surveys*, 17, 169–215. <https://doi.org/10.1214/23-SS144>
- Burrows, A., Sudarsky, D., Sharp, C., Marley, M., Hubbard, W. B., Lunine, J. I., Guillot, T., Saumon, D., & Freedman, R. (1997). Advances in the theory of extrasolar giant planets and brown dwarfs. *arXiv e-Prints*, astro-ph/9706080. <https://doi.org/10.48550/arXiv.astro-ph/9706080>
- Castro-González, A., Demangeon, O. D. S., Lillo-Box, J., Lovis, C., Lavie, B., Adibekyan, V., Acuña, L., Deleuil, M., Aguichine, A., Zapatero Osorio, M. R., Tabernero, H. M., Davout, J., Alibert, Y., Santos, N., Sousa, S. G., Antoniadis-Karnavas, A., Borsa, F., Winn, J. N., Allende Prieto, C., ... Suárez Mascareño, A. (2023). An unusually low-density super-Earth transiting the bright early-type M-dwarf GJ 1018 (TOI-244). *Astronomy & Astrophysics*, 675, A52. <https://doi.org/10.1051/0004-6361/202346550>

- Chabrier, G., & Debras, F. (2021). A new equation of state for dense hydrogen-helium mixtures. II. Taking into account hydrogen-helium interactions. *The Astrophysical Journal*, 917(1), 4. <https://doi.org/10.3847/1538-4357/abfc48>
- Charnay, B., Bézard, B., Baudino, J.-L., Bonnefoy, M., Boccaletti, A., & Galicher, R. (2018). A self-consistent cloud model for brown dwarfs and young giant exoplanets: Comparison with photometric and spectroscopic observations. *The Astrophysical Journal*, 854(2), 172. <https://doi.org/10.3847/1538-4357/aaac7d>
- Chen, H., & Rogers, L. A. (2016). Evolutionary analysis of gaseous sub-Neptune-mass planets with MESA. *The Astrophysical Journal*, 831(2), 180. <https://doi.org/10.3847/0004-637X/831/2/180>
- Dorn, C., Khan, A., Heng, K., Connolly, J. A. D., Alibert, Y., Benz, W., & Tackley, P. (2015). Can we constrain the interior structure of rocky exoplanets from mass and radius measurements? *Astronomy & Astrophysics*, 577, A83. <https://doi.org/10.1051/0004-6361/201424915>
- Egger, J. A., Osborn, H. P., Kubyshkina, D., Mordasini, C., Alibert, Y., Günther, M. N., Lendl, M., Brandeker, A., Heitzmann, A., Leleu, A., Damasso, M., Bonfanti, A., Wilson, T. G., Sousa, S. G., Haldemann, J., Delrez, L., Hooton, M. J., Zingales, T., Luque, R., ... Walton, N. A. (2024). Unveiling the internal structure and formation history of the three planets transiting HIP 29442 (TOI-469) with CHEOPS. *Astronomy & Astrophysics*, 688, A223. <https://doi.org/10.1051/0004-6361/202450472>
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. (2013). emcee: The MCMC Hammer. *Publications of the Astronomical Society of the Pacific*, 125(925), 306. <https://doi.org/10.1086/670067>
- Fortney, J. J., Mordasini, C., Nettelmann, N., Kempton, E. M.-R., Greene, T. P., & Zahnle, K. (2013). A framework for characterizing the atmospheres of low-mass low-density transiting planets. *The Astrophysical Journal*, 775(1), 80. <https://doi.org/10.1088/0004-637X/775/1/80>
- Ginzburg, S., & Chiang, E. (2020). Heavy-metal Jupiters by major mergers: metallicity versus mass for giant planets. *Monthly Notices of the Royal Astronomical Society*, 498(1), 680–688. <https://doi.org/10.1093/mnras/staa2500>
- Haldemann, J., Alibert, Y., Mordasini, C., & Benz, W. (2020). AQUA: a collection of H₂O equations of state for planetary models. *Astronomy & Astrophysics*, 643, A105. <https://doi.org/10.1051/0004-6361/202038367>
- Haldemann, J., Dorn, C., Venturini, J., Alibert, Y., & Benz, W. (2024). BICEPS: An improved characterization model for low- and intermediate-mass exoplanets. *Astronomy & Astrophysics*, 681, A96. <https://doi.org/10.1051/0004-6361/202346965>
- Helled, R., Werner, S., Dorn, C., Guillot, T., Ikoma, M., Ito, Y., Kama, M., Lichtenberg, T., Miguel, Y., Shorttle, O., Tackley, P. J., Valencia, D., & Vazan, A. (2022). Ariel planetary interiors White Paper. *Experimental Astronomy*, 53(2), 323–356. <https://doi.org/10.1007/s10686-021-09739-3>
- Howard, S., & Guillot, T. (2023). Accounting for non-ideal mixing effects in the hydrogen-helium equation of state. *Astronomy & Astrophysics*, 672, L1. <https://doi.org/10.1051/0004-6361/202244851>
- Howard, S., Guillot, T., Bazot, M., Miguel, Y., Stevenson, D. J., Galanti, E., Kaspi, Y., Hubbard, W. B., Militzer, B., Helled, R., Nettelmann, N., Idini, B., & Bolton, S. (2023). Jupiter's interior from Juno: Equation-of-state uncertainties and dilute core extent. *Astronomy & Astrophysics*, 672, A33. <https://doi.org/10.1051/0004-6361/202245625>
- Huang, C., Rice, D. R., & Steffen, J. H. (2022). MAGRATHEA: an open-source spherical

- symmetric planet interior structure code. *Monthly Notices of the Royal Astronomical Society*, 513(4), 5256–5269. <https://doi.org/10.1093/mnras/stac1133>
- Irwin, P. G. J., Dobinson, J., James, A., Toledo, D., Teanby, N. A., Fletcher, L. N., Orton, G. S., & Pérez-Hoyos, S. (2021). Latitudinal variation of methane mole fraction above clouds in Neptune's atmosphere from VLT/MUSE-NFM: Limb-darkening reanalysis. *Icarus*, 357, 114277. <https://doi.org/10.1016/j.icarus.2020.114277>
- Jørgensen, U. G., Amadio, F., Campos Estrada, B., Møller, K. H., Schneider, A. D., Balduin, T., D'Alessandro, A., Symeonidou, E., Helling, C., Nordlund, Å., & Woitke, P. (2024). A grid of self-consistent MSG (MARCS-StaticWeather-GGchem) cool stellar, sub-stellar, and exoplanetary model atmospheres. *Astronomy & Astrophysics*, 690, A127. <https://doi.org/10.1051/0004-6361/202450108>
- Karalidi, T., Marley, M., Fortney, J. J., Morley, C., Saumon, D., Lupu, R., Visscher, C., & Freedman, R. (2021). The Sonora Substellar Atmosphere Models. II. Cholla: A grid of cloud-free, solar metallicity models in chemical disequilibrium for the JWST era. *The Astrophysical Journal*, 923(2), 269. <https://doi.org/10.3847/1538-4357/ac3140>
- Karkoschka, E., & Tomasko, M. G. (2011). The haze and methane distributions on Neptune from HST-STIS spectroscopy. *Icarus*, 211(1), 780–797. <https://doi.org/10.1016/j.icarus.2010.08.013>
- Kempton, E. M.-R., Lessard, M., Malik, M., Rogers, L. A., Futrowsky, K. E., Ih, J., Marounina, N., & Romero-Mirza, C. E. (2023). Where are the water worlds?: Self-consistent models of water-rich exoplanet atmospheres. *The Astrophysical Journal*, 953(1), 57. <https://doi.org/10.3847/1538-4357/ace10d>
- Lacy, B., & Burrows, A. (2023). Self-consistent models of Y dwarf atmospheres with water clouds and disequilibrium chemistry. *The Astrophysical Journal*, 950(1), 8. <https://doi.org/10.3847/1538-4357/acc8cb>
- Linder, E. F., Mordasini, C., Mollière, P., Marleau, G.-D., Malik, M., Quanz, S. P., & Meyer, M. R. (2019). Evolutionary models of cold and low-mass planets: cooling curves, magnitudes, and detectability. *Astronomy & Astrophysics*, 623, A85. <https://doi.org/10.1051/0004-6361/201833873>
- Lopez, E. D., & Fortney, J. J. (2014). Understanding the mass-radius relation for sub-neptunes: Radius as a proxy for composition. *The Astrophysical Journal*, 792(1), 1. <https://doi.org/10.1088/0004-637X/792/1/1>
- Lyon, S. P. (1992). Sesame: The Los Alamos National Laboratory equation of state database. *Los Alamos National Laboratory Report LA-UR-92-3407*.
- Marley, M. S., Fortney, J., Seager, S., & Barman, T. (2007). Atmospheres of extrasolar giant planets. In B. Reipurth, D. Jewitt, & K. Keil (Eds.), *Protostars and Planets V* (p. 733). <https://doi.org/10.48550/arXiv.astro-ph/0602468>
- Marley, M. S., Saumon, D., Visscher, C., Lupu, R., Freedman, R., Morley, C., Fortney, J. J., Seay, C., Smith, A. J. R. W., Teal, D. J., & Wang, R. (2021). The Sonora Brown Dwarf Atmosphere and Evolution Models. I. Model description and application to cloudless atmospheres in rainout chemical equilibrium. *The Astrophysical Journal*, 920(2), 85. <https://doi.org/10.3847/1538-4357/ac141d>
- Mazevet, S., Licari, A., Chabrier, G., & Potekhin, A. Y. (2019). Ab initio based equation of state of dense water for planetary and exoplanetary modeling. *Astronomy & Astrophysics*, 621, A128. <https://doi.org/10.1051/0004-6361/201833963>
- Miguel, Y., Bazot, M., Guillot, T., Howard, S., Galanti, E., Kaspi, Y., Hubbard, W. B., Militzer, B., Helled, R., Atreya, S. K., Connerney, J. E. P., Durante, D., Kulowski, L., Lunine, J. I., Stevenson, D., & Bolton, S. (2022). Jupiter's inhomogeneous envelope. *Astronomy &*

- Astrophysics*, 662, A18. <https://doi.org/10.1051/0004-6361/202243207>
- Miguel, Y., & Vazan, A. (2023). Interior and evolution of the giant planets. *Remote Sensing*, 15(3), 681. <https://doi.org/10.3390/rs15030681>
- Mollière, P., van Boekel, R., Bouwman, J., Henning, Th., Lagage, P.-O., & Min, M. (2017). Observing transiting planets with JWST. Prime targets and their synthetic spectral observations. *Astronomy & Astrophysics*, 600, A10. <https://doi.org/10.1051/0004-6361/201629800>
- Mollière, P., van Boekel, R., Dullemond, C., Henning, Th., & Mordasini, C. (2015). Model atmospheres of irradiated exoplanets: The influence of stellar parameters, metallicity, and the C/O ratio. *The Astrophysical Journal*, 813(1), 47. <https://doi.org/10.1088/0004-637X/813/1/47>
- Morley, C. V., Mukherjee, S., Marley, M. S., Fortney, J. J., Visscher, C., Lupu, R., Gharib-Nezhad, E., Thorngren, D., Freedman, R., & Batalha, N. (2024). The Sonora Substellar Atmosphere Models. III. Diamondback: Atmospheric properties, spectra, and evolution for warm cloudy substellar objects. *The Astrophysical Journal*, 975(1), 59. <https://doi.org/10.3847/1538-4357/ad71d5>
- Mosegaard, K., & Tarantola, A. (1995). Monte Carlo sampling of solutions to inverse problems. *Journal of Geophysical Research: Solid Earth*, 100(B7), 12, 431–412, 447. <https://doi.org/10.1029/94JB03097>
- Mousis, O., Deleuil, M., Aguichine, A., Marcq, E., Naar, J., Aguirre, L. A., Brugger, B., & Gonçalves, T. (2020). Irradiated ocean planets bridge super-Earth and sub-Neptune populations. *The Astrophysical Journal Letters*, 896(2), L22. <https://doi.org/10.3847/2041-8213/ab9530>
- Mukherjee, S., Fortney, J. J., Morley, C. V., Batalha, N. E., Marley, M. S., Karalidi, T., Visscher, C., Lupu, R., Freedman, R., & Gharib-Nezhad, E. (2024). The Sonora Substellar Atmosphere Models. IV. Elf Owl: Atmospheric mixing and chemical disequilibrium with varying metallicity and C/O ratios. *The Astrophysical Journal*, 963(1), 73. <https://doi.org/10.3847/1538-4357/ad18c2>
- Müller, S., & Helled, R. (2021). Synthetic evolution tracks of giant planets. *Monthly Notices of the Royal Astronomical Society*, 507(2), 2094–2102. <https://doi.org/10.1093/mnras/stab2250>
- Otegi, J. F., Dorn, C., Helled, R., Bouchy, F., Haldemann, J., & Alibert, Y. (2020). Impact of the measured parameters of exoplanets on the inferred internal structure. *Astronomy & Astrophysics*, 640, A135. <https://doi.org/10.1051/0004-6361/202038006>
- Owen, J. E., & Wu, Y. (2013). Kepler planets: A tale of evaporation. *The Astrophysical Journal*, 775(2), 105. <https://doi.org/10.1088/0004-637X/775/2/105>
- Paxton, B., Bildsten, L., Dotter, A., Herwig, F., Lesaffre, P., & Timmes, F. (2011). Modules for Experiments in Stellar Astrophysics (MESA). *The Astrophysical Journal Supplement Series*, 192(1), 3. <https://doi.org/10.1088/0067-0049/192/1/3>
- Paxton, B., Cantiello, M., Arras, P., Bildsten, L., Brown, E. F., Dotter, A., Mankovich, C., Montgomery, M. H., Stello, D., Timmes, F. X., & Townsend, R. (2013). Modules for Experiments in Stellar Astrophysics (MESA): Planets, oscillations, rotation, and massive stars. *The Astrophysical Journal Supplement Series*, 208(1), 4. <https://doi.org/10.1088/0067-0049/208/1/4>
- Paxton, B., Marchant, P., Schwab, J., Bauer, E. B., Bildsten, L., Cantiello, M., Dessart, L., Farmer, R., Hu, H., Langer, N., Townsend, R. H. D., Townsley, D. M., & Timmes, F. X. (2015). Modules for Experiments in Stellar Astrophysics (MESA): Binaries, pulsations, and explosions. *The Astrophysical Journal Supplement Series*, 220(1), 15. <https://doi.org/10.1088/0067-0049/220/1/15>

- Paxton, B., Schwab, J., Bauer, E. B., Bildsten, L., Blinnikov, S., Duffell, P., Farmer, R., Goldberg, J. A., Marchant, P., Sorokina, E., Thoul, A., Townsend, R. H. D., & Timmes, F. X. (2018). Modules for Experiments in Stellar Astrophysics (MESA): Convective boundaries, element diffusion, and massive star explosions. *The Astrophysical Journal Supplement Series*, 234(2), 34. <https://doi.org/10.3847/1538-4365/aaa5a8>
- Paxton, B., Smolec, R., Schwab, J., Gautschy, A., Bildsten, L., Cantiello, M., Dotter, A., Farmer, R., Goldberg, J. A., Jermyn, A. S., Kanbur, S. M., Marchant, P., Thoul, A., Townsend, R. H. D., Wolf, W. M., Zhang, M., & Timmes, F. X. (2019). Modules for Experiments in Stellar Astrophysics (MESA): Pulsating variable stars, rotation, convective boundaries, and energy conservation. *The Astrophysical Journal Supplement Series*, 243(1), 10. <https://doi.org/10.3847/1538-4365/ab2241>
- Piaulet, C., Benneke, B., Rubenzahl, R. A., Howard, A. W., Lee, E. J., Thorngren, D., Angus, R., Peterson, M., Schlieder, J. E., Werner, M., Kreidberg, L., Jaouni, T., Crossfield, I. J. M., Ciardi, D. R., Petigura, E. A., Livingston, J., Dressing, C. D., Fulton, B. J., Beichman, C., ... Sinukoff, E. (2021). WASP-107b's density is even lower: A case study for the physics of planetary gas envelope accretion and orbital migration. *The Astronomical Journal*, 161(2), 70. <https://doi.org/10.3847/1538-3881/abcd3c>
- Poser, A. J., Nettelmann, N., & Redmer, R. (2019). The effect of clouds as an additional opacity source on the inferred metallicity of giant exoplanets. *Atmosphere*, 10(11), 664. <https://doi.org/10.3390/atmos10110664>
- Poser, A. J., & Redmer, R. (2024). The effect of cloudy atmospheres on the thermal evolution of warm giant planets from an interior modelling perspective. *Monthly Notices of the Royal Astronomical Society*, 529(3), 2242–2257. <https://doi.org/10.1093/mnras/stae645>
- Rogers, J. G., Owen, J. E., & Schlichting, H. E. (2024). Under the light of a new star: evolution of planetary atmospheres through protoplanetary disc dispersal and boil-off. *Monthly Notices of the Royal Astronomical Society*, 529(3), 2716–2733. <https://doi.org/10.1093/mnras/stae563>
- Rogers, J. G., Schlichting, H. E., & Owen, J. E. (2023). Conclusive evidence for a population of water worlds around M dwarfs remains elusive. *The Astrophysical Journal Letters*, 947(1), L19. <https://doi.org/10.3847/2041-8213/acc86f>
- Roth, A., Parmentier, V., & Hammond, M. (2024). Hot Jupiter diversity and the onset of TiO/VO revealed by a large grid of non-grey global circulation models. *Monthly Notices of the Royal Astronomical Society*, 531(1), 1056–1083. <https://doi.org/10.1093/mnras/stae984>
- Samland, M., Mollière, P., Bonnefoy, M., Maire, A.-L., Cantalloube, F., Cheetham, A. C., Mesa, D., Gratton, R., Biller, B. A., Wahhaj, Z., Bouwman, J., Brandner, W., Melnick, D., Carson, J., Janson, M., Henning, T., Homeier, D., Mordasini, C., Langlois, M., ... Weber, L. (2017). Spectral and atmospheric characterization of 51 Eridani b using VLT/SPHERE. *Astronomy & Astrophysics*, 603, A57. <https://doi.org/10.1051/0004-6361/201629767>
- Saumon, D., Chabrier, G., & van Horn, H. M. (1995). An equation of state for low-mass stars and giant planets. *The Astrophysical Journal Supplement Series*, 99, 713. <https://doi.org/10.1086/192204>
- Schneider, A. D., & Bitsch, B. (2021). How drifting and evaporating pebbles shape giant planets. I. Heavy element content and atmospheric C/O. *Astronomy & Astrophysics*, 654, A71. <https://doi.org/10.1051/0004-6361/202039640>
- Seager, S., Kuchner, M., Hier-Majumder, C. A., & Militzer, B. (2007). Mass-radius relationships for solid exoplanets. *The Astrophysical Journal*, 669(2), 1279–1297. <https://doi.org/10.1086/521346>
- Suissa, G., Chen, J., & Kipping, D. (2018). A HARDCORE model for constraining an exo-

- planet's core size. *Monthly Notices of the Royal Astronomical Society*, 476(2), 2613–2620. <https://doi.org/10.1093/mnras/sty381>
- Teske, J. K., Thorngren, D., Fortney, J. J., Hinkel, N., & Brewer, J. M. (2019). Do metal-rich stars make metal-rich planets? New insights on giant planet formation from host star abundances. *The Astronomical Journal*, 158(6), 239. <https://doi.org/10.3847/1538-3881/ab4f79>
- Thomas, S. W., & Madhusudhan, N. (2016). In hot water: effects of temperature-dependent interiors on the radii of water-rich super-Earths. *Monthly Notices of the Royal Astronomical Society*, 458(2), 1330–1344. <https://doi.org/10.1093/mnras/stw321>
- Thorngren, D. P., Fortney, J. J., Murray-Clay, R. A., & Lopez, E. D. (2016). The mass-metallicity relation for giant planets. *The Astrophysical Journal*, 831(1), 64. <https://doi.org/10.3847/0004-637X/831/1/64>
- Turbet, M., Bolmont, E., Ehrenreich, D., Gratier, P., Leconte, J., Selsis, F., Hara, N., & Lovis, C. (2020). Revised mass-radius relationships for water-rich rocky planets more irradiated than the runaway greenhouse limit. *Astronomy & Astrophysics*, 638, A41. <https://doi.org/10.1051/0004-6361/201937151>
- Unterborn, C. T., Desch, S. J., Haldemann, J., Lorenzo, A., Schulze, J. G., Hinkel, N. R., & Panero, W. R. (2023). The nominal ranges of rocky planet masses, radii, surface gravities, and bulk densities. *The Astrophysical Journal*, 944(1), 42. <https://doi.org/10.3847/1538-4357/acaa3b>
- Vivien, H. G., Aguichine, A., Mousis, O., Deleuil, M., & Marcq, E. (2022). On the stability of low-mass planets with supercritical hydrospheres. *The Astrophysical Journal*, 931(2), 143. <https://doi.org/10.3847/1538-4357/ac66e2>
- Wang, H. S., Liu, F., Ireland, T. R., Brasser, R., Yong, D., & Lineweaver, C. H. (2019). Enhanced constraints on the interior composition and structure of terrestrial exoplanets. *Monthly Notices of the Royal Astronomical Society*, 482(2), 2222–2233. <https://doi.org/10.1093/mnras/sty2749>
- Zeng, L., Jacobsen, S. B., Sasselov, D. D., Petaev, M. I., Vanderburg, A., Lopez-Morales, M., Perez-Mercader, J., Mattsson, T. R., Li, G., Heising, M. Z., Bonomo, A. S., Damasso, M., Berger, T. A., Cao, H., Levi, A., & Wordsworth, R. D. (2019). Growth model interpretation of planet size distribution. *Proceedings of the National Academy of Science*, 116(20), 9723–9728. <https://doi.org/10.1073/pnas.1812905116>
- Zeng, L., Sasselov, D. D., & Jacobsen, S. B. (2016). Mass-radius relation for rocky planets based on PREM. *The Astrophysical Journal*, 819(2), 127. <https://doi.org/10.3847/0004-637X/819/2/127>