

Limace.jl: A Julia package to compute hydromagnetic modes in spherical domains

Felix Gerick ^[]^{1,2,3}

1 Paris Institute of Earth Physics (IPGP) ROR 2 National Centre for Space Studies (CNES), France ROR 3 Royal Observatory of Belgium, Belgium ROR

Summary

Hydromagnetic modes in spherical domains are relevant to the liquid cores of planets, moons or stars, as well as rotating fluid dynamics experiments. These modes are solutions to the linearized rotating magnetohydrodynamic equations that govern electrically conducting fluids under rapid rotation. Limace.jl is a package written in the Julia programming language (Bezanson et al., 2017), based on Galerkin projections of the governing equations onto trial vectors of the velocity and magnetic field. It aims to facilitate the calculation of modes in flexible setups with a high-level interface, whilst remaining computationally performant enough to tackle relevant physical parameters.

Statement of need

The study of hydromagnetic modes is relevant in particular to Earth's liquid core. Despite having been theoretically predicted a long time ago (Braginsky, 1970; Hide, 1966; Malkus, 1967), recent advances in numerical modelling and new observational evidence in geomagnetic data have reignited interest in these modes (Gerick et al., 2021; Gerick & Livermore, 2024; Gillet et al., 2022; Luo et al., 2022; Luo & Jackson, 2022; Triana et al., 2022). It is therefore relevant to the geophysical and astrophysical fluid dynamics community to have access to a code that models these modes. The only open-source models to compute hydromagnetic modes in relevant parameters for planetary cores is Kore (Triana et al., 2021), a spectral code based on ultraspherical polynomials in axisymmetric setups written in Python. Although Kore is limited to axisymmetric geometries, the implementation requires substantial entry barrier for scientists to model these modes. In addition to Kore, Limace.jl tries to lower this entry barrier, by providing a generic open source model with a very simple high-level API and modern online documentation with practical examples.

Despite having a high-level interface, Limace.jl can be used to solve complex and geophysically relevant problems. A unique feature of Limace.jl is the support of complex (non-axisymmetric) background magnetic fields and flows over which the modes evolve. The code has been developed from the beginning to leave assumptions of symmetry up to the user. The model code base is tested against mode solutions from the scientific literature to ensure its correctness.

Theoretical background and implementation details

In order to compute modal solutions, we consider the linearized momentum equation of the incompressible fluid and the linearized induction equation (Gerick & Livermore, 2024; Ivers &

DOI: 10.21105/joss.07593

Software

- Review 2
- Repository ¹
- Archive I^a

Editor: Patrick Diehl ♂ ◎ Reviewers:

- @milankl
 - Øjakelangham

Submitted: 07 October 2024 Published: 18 June 2025

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

Gerick. (2025). Limace.jl: A Julia package to compute hydromagnetic modes in spherical domains. *Journal of Open Source Software*, 10(110), 1 7593. https://doi.org/10.21105/joss.07593.



Phillips, 2008)

$$\begin{split} \lambda \mathbf{u} &= -\left(\boldsymbol{\nabla}\times\mathbf{u}\right)\times\mathbf{U}_0 - \left(\boldsymbol{\nabla}\times\mathbf{U}_0\right)\times\mathbf{u} - 2\boldsymbol{\Omega}\times\mathbf{u} - \frac{1}{\rho}\nabla p \\ &+ \frac{1}{\rho\mu_0}\left(\left(\boldsymbol{\nabla}\times\mathbf{b}\right)\times\mathbf{B}_0 + \left(\boldsymbol{\nabla}\times\mathbf{B}_0\right)\times\mathbf{b}\right) + \nu\boldsymbol{\nabla}^2\mathbf{u}, \\ \lambda \mathbf{b} &= \boldsymbol{\nabla}\times\left(\mathbf{U}_0\times\mathbf{b}\right) + \boldsymbol{\nabla}\times\left(\mathbf{u}\times\mathbf{B}_0\right) + \eta\boldsymbol{\nabla}^2\mathbf{b}. \end{split}$$

with **u** the velocity perturbation, \mathbf{U}_0 the steady background velocity, **b** the magnetic field perturbation, \mathbf{B}_0 the background magnetic field, $\boldsymbol{\Omega}$ the rotation axis, ρ the fluid density, p the reduced hydrodynamic pressure, μ_0 the magnetic permeability of free space, ν the kinematic viscosity, η the magnetic diffusivity, and $\lambda = -\sigma + i\omega$, with σ the damping rate and ω the frequency of the oscillatory perturbation to the steady background.

To discretize the linearized equations, they are projected onto trial vectors (Galerkin vector bases) \mathbf{u}_i and \mathbf{b}_i for the velocity and magnetic field, respectively. Due to the divergence free condition on the velocity and magnetic field, i.e. the flow is incompressible and no magnetic monopoles exist, it is convenient to decompose the fields into poloidal and toroidal components, so that

$$\mathbf{u} = \sum_{i} \alpha_{i} \mathbf{u}_{i} = \sum_{l,m,n} \alpha_{lmn}^{P} \mathbf{P}_{lmn} + \sum_{l,m,n} \alpha_{lmn}^{Q} \mathbf{Q}_{lmn},$$
$$\mathbf{b} = \sum_{i} \beta_{i} \mathbf{b}_{i} = \sum_{l,m,n} \beta_{lmn}^{S} \mathbf{S}_{lmn} + \sum_{l,m,n} \beta_{lmn}^{T} \mathbf{T}_{lmn},$$

with $\alpha_i, \beta_i \in \mathbb{C}$. The respective poloidal \mathbf{P}_{lmn} and toroidal \mathbf{Q}_{lmn} basis vectors in spherical coordinates (r, θ, ϕ) are

$$\begin{aligned} \mathbf{P}_{lmn} &= \boldsymbol{\nabla} \times \boldsymbol{\nabla} \times P_{ln}(r) Y_l^m(\theta, \phi) \mathbf{r}, \\ \mathbf{Q}_{lmn} &= \boldsymbol{\nabla} \times Q_{ln}(r) Y_l^m(\theta, \phi) \mathbf{r}, \end{aligned}$$

and analogously for S_{lmn} and T_{lmn} . Here, $Y_l^m(\theta, \phi)$ is the (fully normalized) spherical harmonic of degree l and order m. The boundary conditions (or regularity condition at r = 0) are imposed on the scalar functions P, S, Q, T. The scalar functions can be chosen to have optimal properties, i.e. the resulting basis is orthogonal w.r.t a given inner product (Chen et al., 2018; Gerick & Livermore, 2024; Livermore, 2010). Limace.jl provides several optimal bases that satisfy relevant boundary conditions, such as non-penetration and no-slip for the velocity and perfectly conducting and insulating exteriors for the magnetic field.

We need to consider all combinations of poloidal and toroidal vector combinations in the projection of the forces. This leads to several coupling terms, especially for the Lorentz force, advection and induction terms. The integrals of these coupling terms over the spherical surfaces are computed through the Adam-Gaunt and Elsasser variables (James, 1973), which are calculated from Wigner symbols (available in Julia through WignerSymbols.jl, based on Johansson & Forssén (2016)). The remaining integration in radial direction is done using Gauss-Legendre quadratures, available through FastGaussQuadrature.jl. The exact modelled equations are outlined in Gerick & Livermore (2024), based on the work of Ivers & Phillips (2008).

From the projected equations, the problem reduces to a generalized eigenproblem

$$\lambda \mathbf{A}\mathbf{x} = \mathbf{B}\mathbf{x}$$

that is solved numerically. Here, the eigenvector \mathbf{x} contains the coefficients α_i (and β_i). The matrix \mathbf{B} is generally not symmetric/Hermitian, but \mathbf{A} can be the unit matrix, symmetric tridiagonal or symmetric, depending on the chosen bases.

For small problem sizes, the eigenproblem can be solved using dense methods, e.g. using the standard library function LinearAlgebra.eigen. For larger problem sizes it is only feasible



to compute a few eigensolutions of the sparse system. To this end, a shift-invert spectral transform method is provided, based on the sparse LU factorization from UMFPACK (Davis, 2004) and the partial Schur decomposition implemented in ArnoldiMethod.jl (Stoppels & Nyman, n.d.).

For postprocessing, Limace.jl uses a fast spherical harmonic transform implemented in the SHTns library (Schaeffer, 2013), and available in Julia through SHTns.jl. It is used to transform the spectral coefficients to vector fields evaluated on a spatial grid. Limace.jl does not provide any plotting routines, but some examples are given, leaving the choice of plotting library up to the user.

Acknowledgements

I have received funding from the European Research Council (ERC) GRACEFUL Synergy Grant No. 855677. This project has been funded by ESA in the framework of EO Science for Society, through contract 4000127193/19/NL/IA (SWARM + 4D Deep Earth: Core).

I thank Phil Livermore for the key contributions in the theoretical development of the model.

References

- Bezanson, J., Edelman, A., Karpinski, S., & Shah, V. B. (2017). Julia: A Fresh Approach to Numerical Computing. SIAM Review, 59(1), 65–98. https://doi.org/10.1137/141000671
- Braginsky, S. I. (1970). Torsional magnetohydrodynamics vibrations in the Earth's core and variations in day length. *Geomagnetism and Aeronomy*, *10*, 3–12.
- Chen, L., Herreman, W., Li, K., Livermore, P. W., Luo, J. W., & Jackson, A. (2018). The optimal kinematic dynamo driven by steady flows in a sphere. *Journal of Fluid Mechanics*, 839, 1–32. https://doi.org/10.1017/jfm.2017.924
- Davis, T. A. (2004). Algorithm 832: UMFPACK V4.3—an unsymmetric-pattern multifrontal method. ACM Transactions on Mathematical Software, 30(2), 196–199. https://doi.org/ 10.1145/992200.992206
- Gerick, F., Jault, D., & Noir, J. (2021). Fast Quasi-Geostrophic Magneto-Coriolis Modes in the Earth's Core. Geophysical Research Letters, 48(4), e2020GL090803. https://doi.org/ 10.1029/2020GL090803
- Gerick, F., & Livermore, P. W. (2024). Interannual Magneto-Coriolis modes and their sensitivity on the magnetic field within the Earth's core. *Proceedings of the Royal Society* A: Mathematical, Physical and Engineering Sciences, 480(2299), 20240184. https://doi. org/10.1098/rspa.2024.0184
- Gillet, N., Gerick, F., Jault, D., Schwaiger, T., Aubert, J., & Istas, M. (2022). Satellite magnetic data reveal interannual waves in Earth's core. *Proceedings of the National Academy of Sciences*, 119(13), e2115258119. https://doi.org/10.1073/pnas.2115258119
- Hide, R. (1966). Free hydromagnetic oscillations of the Earth's core and the theory of the geomagnetic secular variation. *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences, 259*(1107), 615–647. https: //doi.org/10.1098/rsta.1966.0026
- Ivers, D. J., & Phillips, C. G. (2008). Scalar and vector spherical harmonic spectral equations of rotating magnetohydrodynamics. *Geophysical Journal International*, 175(3), 955–974. https://doi.org/10.1111/j.1365-246X.2008.03944.x
- James, R. W. (1973). The Adams and Elsasser Dynamo Integrals. Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences, 331(1587), 469–478.



https://www.jstor.org/stable/78350

- Johansson, H. T., & Forssén, C. (2016). Fast and Accurate Evaluation of Wigner 3*j*, 6*j*, and 9*j* Symbols Using Prime Factorization and Multiword Integer Arithmetic. SIAM Journal on Scientific Computing, 38(1), A376–A384. https://doi.org/10.1137/15M1021908
- Livermore, P. W. (2010). Galerkin orthogonal polynomials. *Journal of Computational Physics*, 229(6), 2046–2060. https://doi.org/10.1016/j.jcp.2009.11.022
- Luo, J., & Jackson, A. (2022). Waves in the Earth's core. I. Mildly diffusive torsional oscillations. Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences, 478(2259), 20210982. https://doi.org/10.1098/rspa.2021.0982
- Luo, J., Marti, P., & Jackson, A. (2022). Waves in the Earth's core. II. Magneto-Coriolis modes. Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences, 478(2261), 20220108. https://doi.org/10.1098/rspa.2022.0108
- Malkus, W. V. R. (1967). Hydromagnetic planetary waves. *Journal of Fluid Mechanics*, 28(4), 793–802. https://doi.org/10.1017/S0022112067002447
- Schaeffer, N. (2013). Efficient spherical harmonic transforms aimed at pseudospectral numerical simulations. Geochemistry, Geophysics, Geosystems, 14(3), 751–758. https://doi.org/10. 1002/ggge.20071
- Stoppels, H. T., & Nyman, L. (n.d.). ArnoldiMethod.jl: Arnoldi Method with Krylov-Schur restart, natively in Julia. https://github.com/JuliaLinearAlgebra/ArnoldiMethod.jl
- Triana, S. A., Dumberry, M., Cébron, D., Vidal, J., Trinh, A., Gerick, F., & Rekier, J. (2022). Core Eigenmodes and their Impact on the Earth's Rotation. *Surveys in Geophysics*, 43(1), 107–148. https://doi.org/10.1007/s10712-021-09668-y
- Triana, S. A., Trinh, A., Rekier, J., Zhu, P., & Dehant, V. (2021). The Viscous and Ohmic Damping of the Earth's Free Core Nutation. *Journal of Geophysical Research: Solid Earth*, 126(4), e2020JB021042. https://doi.org/10.1029/2020JB021042