

BatchPlanet: Batch access and processing of PlanetScope imagery for spatiotemporal analysis in R

Yiluan Song ^{1,2}, Alexandre R. Bevington ³, Kenneth N. Reid ⁴, Jiali Zhu ², Yi Liu ², and Kai Zhu ^{2,1}

¹ Michigan Institute for Data and AI in Society, University of Michigan, Ann Arbor, MI, USA ² School for Environment and Sustainability and Institute for Global Change Biology, University of Michigan, Ann Arbor, MI, USA ³ Department of Geography, Earth and Environmental Sciences, University of Northern British Columbia, Prince George, BC, Canada ⁴ Honorary Fellow - University of Stirling, Scotland, UK

DOI: [10.21105/joss.09268](https://doi.org/10.21105/joss.09268)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [William Gearty](#) 

Reviewers:

- [@ajpelu](#)
- [@QuinnAsena](#)

Submitted: 25 July 2025

Published: 16 March 2026

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

The *BatchPlanet* R package offers a reproducible and scalable workflow for accessing and processing PlanetScope satellite imagery. It is designed for environmental researchers to efficiently perform spatiotemporal analysis on high-resolution remote sensing data. The package streamlines imagery ordering and downloading with the PlanetScope API, retrieval and cleaning of pixel-level time series, calculation of vegetation indices like the Enhanced Vegetation Index (EVI), and calculation of start/end of season metrics. This tool is particularly valuable for research involving large volumes of imagery across multiple sites and extended time periods.

Statement of Need

PlanetScope imagery provides global, high-resolution (~3-meter), near-daily data, making it valuable for scientific research and monitoring of phenology ([Moon et al., 2021](#)), land use change, disaster impacts, and more. While the Planet API facilitates access ([Team, 2017](#)), using this data remains challenging due to complex API interactions, limits on large-volume data downloads, and non-trivial processing workflows, which can hinder reproducibility.

Existing options include the official [Planet Python SDK](#), cloud-based platforms like [Sentinel Hub](#) and [Google Earth Engine \(GEE\)](#), and the R package *planetR* ([Bevington et al., 2019](#)). However, the Python and JavaScript used by the first three platforms may be less familiar to R users. Cloud platforms also restrict user control over processing environments and local data storage. Furthermore, existing tools typically require users to write extensive custom scripts for batch downloading and processing across multiple sites, a process often limited by computing and storage resources. *BatchPlanet* addresses these gaps by providing an R-native tool for batch access and processing of PlanetScope imagery (Table 1). Its streamlined, parallelized functions are designed for scalability, transparency, and reproducibility in scientific data pipelines. This package has supported peer-reviewed research in predicting reproductive phenology in wind-pollinated trees ([Song et al., 2025](#)).

Table 1: Comparison of BatchPlanet with existing tools for PlanetScope data access and processing.

Feature / Tool	Planet Python SDK	Sentinel Hub	Google Earth Engine (GEE)	planetR (Bevington)	BatchPlanet
Primary Language	Python	Python	JavaScript / Python	R	R
Processing Environment	Local/Cloud	Cloud	Cloud	Local	Local
Data Control & Reproducibility	High	Moderate	Low	High	High
Batch Processing	Via scripting/CLI	Supported for enterprise users	Via scripting	Via scripting	Streamlined
Time Series Analysis	Not supported	Limited	Supported	Not supported	Supported
Interactive Visualization	Not supported	Limited	Supported	Not supported	Supported

Key Features

The package is tailored for researchers and practitioners who:

- Conduct time series analyses across spatially dispersed monitoring sites.
- Work primarily in R and seek alternatives to Python-based tools.
- Prioritize reproducibility in remote sensing workflows.
- Use high-performance computing (HPC) infrastructure.
- Need interactive visualization of PlanetScope imagery and processed data products.

A major hurdle in using PlanetScope data is the volume of data and the risk of hitting Planet API rate limits during mass downloads. This could happen, for example, when a phenological study requires years of time series over multiple locations. BatchPlanet provides solutions for streamlined batch ordering and downloading. It searches for images by month and automatically splits large requests into smaller orders, ensuring complete data retrieval without hitting API rate limits. It enables users to specify multiple, spatially distant sites, allowing for efficient, parallelized downloading of only the relevant imagery, minimizing unnecessary data volume.

In addition to ordering and downloading, BatchPlanet facilitates the entire R-native workflow for PlanetScope imagery processing, with a focus on temporal analysis. These include functions to retrieve pixel-level time series data, clean reflectance time series, calculate the Normalized Difference Vegetation Index (NDVI) and Enhanced Vegetation Index (EVI) (Huete et al., 2002), and compute start/end of season metrics (the day-of-year when the specified index first crosses specified thresholds) (Moon et al., 2021). Apart from the streamlined batch processing functions, BatchPlanet provides individual functions for key steps of the workflow, allowing users to customize their data processing pipelines. BatchPlanet also enables interactive visualization of true color images and EVI time series (Fig. 1, 2).

These batch functions significantly accelerate the process. For example, images for an approximately 9 km² area over one month were downloaded in 6.7 seconds. The total

downloading time for a year's worth of images is comparable, due to parallelization across months. Time series retrieval (reflectances, quality masks, and metadata) for 100 coordinates from one month of downloaded images took only 20.9 seconds, which can be parallelized across sites and coordinate groups.

Select Site:

AT ▼

Brightness

0 7.5 10

Select Date and Time:

2025-05-11 17:40:57 2025-05-22 17:30:59




Figure 1: A screenshot of the interactive PlanetScope imagery viewer in the BatchPlanet package, showing a true color image in part of Austin, USA, captured on May 11, 2025.




Figure 2: Enhanced Vegetation Index (EVI) for three trees in San Joaquin Experimental range (SJER) NEON site with start/end of season metrics annotated, calculated and visualized using the BatchPlanet package. Points are EVI values calculated from PlanetScope reflectances at the coordinates of the trees of interest, summarized with smoothed lines. Green shades indicate the periods from minimum EVI in the winter to maximum EVI in the summer. Sets of three vertical green lines are the time points when smoothed EVI crosses 30%, 40%, and 50 % (from left to right) of the range between minimum and maximum EVI, which can serve as possible start of season metrics.

Example Usage

Install package in R and load package.

```
install.packages('BatchPlanet',
  repos = c('https://yiluansong.r-universe.dev',
    'https://cloud.r-project.org'))
library(BatchPlanet)
```

To facilitate adoption and ensure reproducibility, BatchPlanet provides three vignettes that demonstrate its full feature set: [Getting started workflow](#), [Customization and advanced usage](#), and [Miscellaneous time series processing tools](#). We highlight some key functions below.

Read example coordinates.

```
df_coordinates <- read.csv(
  system.file("extdata/NEON/example_neon_coordinates.csv",
    package = "BatchPlanet")
)
```

```
visualize_coordinates(df_coordinates)
```

Set download parameters and data directory.

```
setting <- set_planetscope_parameters(
  api_key = set_api_key(),
  item_name = "PSScene",
  asset = "ortho_analytic_4b_sr",
  product_bundle = "analytic_sr_udm2",
```

```
cloud_lim = 0.3,  
harmonized = TRUE  
)  
  
download_sample_data()  
dir_data <- "sample-data"  
dir_data_NEON <- file.path(dir_data, "NEON")
```

Order and download imagery.

Note: Before proceeding to downloading, users should inspect their [Planet account](#) to confirm that all orders reached a “success” status. Failed orders will result in errors during downloading. Refer to the [package vignette](#) for troubleshooting tips when orders fail.

```
order_planetscope_imagery_batch(  
  dir = dir_data_NEON,  
  df_coordinates = df_coordinates,  
  v_site = c("HARV", "SJER"),  
  v_year = 2024,  
  setting = setting  
)
```

```
download_planetscope_imagery_batch(  
  dir = dir_data_NEON,  
  setting = setting,  
  num_cores = 12  
)
```

```
visualize_true_color_imagery_batch(  
  dir = dir_data_NEON,  
  df_coordinates = df_coordinates  
)
```

Retrieve time series at coordinates of interest.

```
retrieve_planetscope_time_series_batch(  
  dir = dir_data_NEON,  
  df_coordinates = df_coordinates,  
  num_cores = 12  
)
```

```
df_ts <- read_data_product(  
  dir = dir_data_NEON,  
  product_type = "ts"  
)
```

```
visualize_time_series(  
  df_ts = df_ts,  
  var = "green",  
  ylab = "Green reflectance",  
  facet_var = "site",  
  smooth = F)
```

Clean time series and calculate NDVI and EVI.

```
clean_planetscope_time_series_batch(  
  dir = dir_data_NEON,  
  num_cores = 3,
```

```
calculate_index = c("ndvi", "evi"),
filter_range = list(ndvi = c(-1, 1), evi = c(0, 1))
)

df_clean <- read_data_product(
  dir = dir_data_NEON,
  product_type = "clean"
)

visualize_time_series(
  df_ts = df_clean,
  var = "evi",
  ylab = "EVI",
  facet_var = "site",
  smooth = T
)

Calculate start and end of season metrics.

df_thres <- set_thresholds(
  thres_up = c(0.3, 0.4, 0.5),
  thres_down = NULL
)

calculate_season_metrics_batch(
  dir = dir_data_NEON,
  v_site = "SJER",
  v_group = "Quercus",
  df_thres = df_thres,
  var_index = "evi",
  num_cores = 12
)

v_id <- c("NEON.PLA.D17.SJER.06001",
         "NEON.PLA.D17.SJER.06337",
         "NEON.PLA.D17.SJER.06310")

df_doy <- read_data_product(
  dir = dir_data_NEON,
  product_type = "doy"
)

df_doy_sample <- df_doy[df_doy$id %in% v_id, ]
df_evi <- read_data_product(
  dir = dir_data_NEON,
  product_type = "clean"
)

df_evi_sample <- df_evi[df_evi$id %in% v_id, ]
visualize_time_series(
  df_ts = df_evi_sample,
  df_doy = df_doy_sample,
  var = "evi",
  ylab = "EVI",
  facet_var = "id",
  smooth = T
)
```

Acknowledgements

Yiluan Song was supported by the Eric and Wendy Schmidt AI in Science Postdoctoral Fellowship, a Schmidt Sciences program. Kai Zhu and Yiluan Song were supported by the National Science Foundation [grant numbers 2306198 (CAREER)]. We thank the Planet team for providing access to their API.

References

- Bevington, A., Tobias, M., Nelsen, T., & Stockwell, E. (2019). *planetR*. <https://github.com/bevingtona/planetR>
- Huete, A., Didan, K., Miura, T., Rodriguez, E. P., Gao, X., & Ferreira, L. G. (2002). Overview of the radiometric and biophysical performance of the MODIS vegetation indices. *Remote Sensing of Environment*, 83(1-2), 195–213. [https://doi.org/10.1016/S0034-4257\(02\)00096-2](https://doi.org/10.1016/S0034-4257(02)00096-2)
- Moon, M., Richardson, A. D., & Friedl, M. A. (2021). Multiscale assessment of land surface phenology from harmonized landsat 8 and sentinel-2, PlanetScope, and PhenoCam imagery. *Remote Sensing of Environment*, 266, 112716. <https://doi.org/10.1016/j.rse.2021.112716>
- Song, Y., Katz, D. S., Zhu, Z., Beaulieu, C., & Zhu, K. (2025). Predicting reproductive phenology of wind-pollinated trees via PlanetScope time series. *Science of Remote Sensing*, 100205. <https://doi.org/10.1016/j.srs.2025.100205>
- Team, P. (2017). *Planet application program interface: In space for life on earth*. <https://api.planet.com>.