

# ThRasE: A QGIS Plugin for Manual Post-Classification Correction and Quality Assurance of Thematic Maps

Xavier C. Llano  <sup>1\*</sup>

<sup>1</sup> Forest and Carbon Monitoring System (SMBYC), IDEAM - Colombia \* These authors contributed equally.

DOI: [10.21105/joss.09634](https://doi.org/10.21105/joss.09634)

## Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

---

Editor: [James Gaboardi](#) 

## Reviewers:

- [@giswqs](#)
- [@Narayana-Rao](#)

Submitted: 07 October 2025

Published: 06 March 2026

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

## Summary

ThRasE is an open-source QGIS plugin designed for systematic review, manual correction, and quality assurance of thematic rasters. It offers synchronized multi-panel views for side-by-side and overlap comparisons with reference data. It utilizes a recode pixel table to modify multiple classes simultaneously in one operation with various editing tool options. An integrated modification tracking system automatically records all changes, providing spatial feedback, transparency, and reproducibility. Systematic tile-based navigation ensures complete spatial coverage during inspection, while session persistence enables seamless multi-day workflows.

## Statement of need

Thematic maps serve as indispensable analytical tools that simplify complex, continuous spatial data into visually accessible representations for a wide range of disciplines and applications. However, mapping workflows inevitably introduce errors, making quality assurance essential for reliable spatial data and informed decision-making (Foody, 2001). Because maps are models of reality, they inevitably contain errors introduced by projection, compilation, and symbolization (Lightfoot & Butler, 1987; Maling, 2013). Many issues originate in the production process itself. For example, satellite-based classifications (e.g., supervised or unsupervised workflows) can introduce errors of omission or commission, scale mismatches, or temporal misalignments with ground conditions (Schowengerdt, 2007). These limitations do not undermine the value of thematic maps; rather, they highlight the importance of rigorous quality assurance. Best practices include conducting formal accuracy assessments (Congalton & Green, 2019), transparently reporting error and uncertainty (MacEachren, 1992), and, where appropriate, applying targeted post-classification corrections to address misclassifications (Manandhar et al., 2009).

The value of a thematic map ultimately depends on its quality, which is primarily driven by the original data sources and classification methodologies. However, post-classification correction serves as a widely adopted final step to refine misclassifications and improve outcomes. A common approach, particularly for remote sensing-based maps, involves integrating ancillary data and knowledge-based rules to resolve misclassifications, reduce commission errors, and enhance overall accuracy (Manandhar et al., 2009; Thakkar et al., 2017). These corrections often rely on manual editing or semi-automated techniques guided by expert knowledge and reference data. This is precisely the context in which ThRasE was developed.

Several tools offer raster editing and reclassification capabilities, including commercial packages such as ERDAS IMAGINE and ENVI with raster reclassification and pixel editing, and ArcGIS with its interactive pixel value manipulation tool. Open-source alternatives like GRASS GIS provide command-line reclassification, and the QGIS plugin Serval enables basic pixel-level

raster modifications. While these tools support raster editing, none integrates synchronized multi-view comparison, systematic tile-based navigation, modification tracking, multiple editing tools, a recode table, and session persistence within a unified workflow.

In Colombia, the Forest and Carbon Monitoring System (SMBByC) of the Institute of Hydrology, Meteorology and Environmental Studies (IDEAM) is responsible for measuring and ensuring the accuracy of the official national forest figures. As part of the SMBByC, we identified the need not only for review and editing capabilities but also for a dedicated tool to enable systematic and rigorous quality control of final products, thereby guaranteeing reliable results (Galindo et al., 2014). To address this gap, we developed ThRasE, an open-source QGIS plugin that provides flexible editing, visual inspection, and auditable quality assurance workflows.

ThRasE offers a suite of editing tools, a modification tracking registry, systematic visual inspection features, and comparison capabilities with reference data. By incorporating ancillary data and knowledge-based rules, ThRasE enables expert-guided manual editing, facilitating targeted corrections of misclassifications and fine-tuning of class assignments. ThRasE has been used for multiple purposes worldwide, including manual post-processing to correct land-use/land-cover misclassifications via photo-interpretation of satellite image mosaics (M. J. Rayner, 2022; Vallet et al., 2024); knowledge-based manual correction of misclassifications and residual errors (Bladh, 2024; M. Rayner, Balzter, et al., 2021; Senterre et al., 2023); manual classification/reclassification of pixels for land-cover reconstruction to reconcile multi-date imagery (Gunawan et al., 2023); annotation of historical aerial imagery for model calibration and fine-tuning (Eyster et al., 2024); as a validation/visual QA step (Hariyanto et al., 2024); refinement of agricultural maps (Gandharum et al., 2025; M. Rayner, Whelan, et al., 2021); among other applications (Dupuy et al., 2024; Hariyanto et al., 2022; Nunes Machado et al., 2024; Queiroga, 2020). Furthermore, ThRasE is an official component of our Digital Image Processing Protocol for Quantifying Deforestation in Colombia (Galindo et al., 2014).

## Key features

### Multi-View Configuration

ThRasE is designed to prioritize visual comparison. To that end, analysts can arrange multiple synchronized panels in a grid, with each panel blending up to three layers (for example, the target thematic raster, a reference product, and ancillary context) with independent opacity controls. Creating several panels with different layer combinations and opacity settings enables overlap inspection and side-by-side comparison, making it easier to spot boundary errors, edge artifacts, and local inconsistencies that are often missed when toggling a single map. This layout shortens the loop between detection and correction during post-classification editing and quality assurance.




Figure 1: Multi-view panels and their layer configuration toolbar with controls for visual comparison.

## Editing and Reclassification Tools

ThRasE provides pixel-level editing and bulk class recoding for fast, large-scale corrections. The Recode Pixel Table lets analysts define class-to-class mappings and apply them across the thematic map, enabling multiple classes to be updated in a single operation. Each view offers an editing toolset with options for pixel, line, polygon, and freehand drawing, supported by independent undo and redo actions per tool and per view. Analysts can initiate edits from any panel while targeting the thematic map, preserving full visual context during correction and facilitating precise, context-aware refinements.




Figure 2: Editing toolbar featuring pixel, line, polygon, and freehand drawing tools.

## Modification Tracking System

ThRasE includes a modification tracking system that records all edits to the thematic map during editing sessions. Every pixel change is recorded with its original value and timestamp, enabling a complete edit history and strong traceability for quality assurance. Modified areas can be highlighted on the canvas to provide immediate spatial feedback and help reveal patterns or missed regions during the editing process. By maintaining a comprehensive record of edits, the system supports quality assurance workflows where transparency and reproducibility are essential, allowing teams to verify corrections, trace decisions, and export change logs for documentation or peer review.





Figure 3: Modification tracking registry showing edit history with timestamps and visual highlighting of modified areas on the canvas.

## Systematic Inspection Navigator

Systematic inspection is fundamental to quality assurance in thematic mapping. ThRasE provides a tile-based navigation tool that divides the area of interest within the thematic map into manageable units and sequences through tiles in a defined pattern to guarantee full spatial coverage. The navigation tool offers multiple configuration options, allowing teams to adapt their review strategy to data characteristics or geographic features. This structured approach transforms what could be an unstructured visual inspection into a methodical process, reducing the risk of missed areas and supporting complete spatial coverage during editing and quality control operations.



**Figure 4:** Tile-based navigation interface dividing the study area into manageable units for systematic inspection with configurable patterns.

## Session Persistence

ThRasE supports session persistence through save and restore functionality that captures the complete workspace state. This capability proves essential for multi-session workflows where analysts need to maintain consistency across editing or review periods. Teams can share standardized configurations to ensure uniform inspection methodology across multiple operators, while individual analysts can resume work at the exact tile and visual context where they paused, preserving both spatial progress and workspace setup.

## Acknowledgements

This project was fully funded by the Forest and Carbon Monitoring System (SMByC) of the Institute of Hydrology, Meteorology and Environmental Studies (IDEAM), Colombia. We sincerely thank the SMByC PDI team for their support in testing, product verification, and providing invaluable feedback throughout the development process.

## References

- Bladh, E. (2024). *Urban Tree Mapping using airborne LiDAR: Analysing vegetation changes between 2010, 2017, and 2022 A Gothenburg case study* [Master's thesis, University of Gothenburg]. <https://hdl.handle.net/2077/81972>
- Congalton, R. G., & Green, K. (2019). *Assessing the Accuracy of Remotely Sensed Data: Principles and Practices*. CRC Press. <https://doi.org/10.1201/9781420048568>
- Dupuy, S., Lelong, C., Gaetano, R., & Villers, A. (2024). *BIOTAMAYA. Rapport méthodologique pour la production de carte d'occupation du sol-Mayotte en 2023*.

- CIRAD-ES-UMR TETIS. <https://agritrop.cirad.fr/608783/>
- Eyster, H. N., Chan, K. M., Fletcher, M. E., & Beckage, B. (2024). Space-for-time substitutions exaggerate urban bird–habitat ecological relationships. *Journal of Animal Ecology*, 93(12), 1854–1867. <https://doi.org/10.1111/1365-2656.14194>
- Foody, G. M. (2001). GIS: The accuracy of spatial data revisited. *Progress in Physical Geography*, 25(3), 389–398. <https://doi.org/10.1177/030913330102500306>
- Galindo, G., Espejo, O. J., Rubiano, J. C., Vergara, L. K., & Cabrera, E. (2014). *Protocolo de procesamiento digital de imágenes para la cuantificación de la deforestación en Colombia. V 2.0*. Instituto de Hidrología, Meteorología y Estudios Ambientales – IDEAM. Bogotá D.C., Colombia. [https://www.ideam.gov.co/sites/default/files/archivos/protocolo\\_de\\_pdi\\_para\\_la\\_cuantificacion\\_de\\_la\\_deforestacion\\_en\\_colombia\\_v2\\_1.pdf](https://www.ideam.gov.co/sites/default/files/archivos/protocolo_de_pdi_para_la_cuantificacion_de_la_deforestacion_en_colombia_v2_1.pdf)
- Gandharum, L., Hartono, D. M., Sadmono, H., Sanjaya, H., Sumargana, L., Kusumawardhani, A. D., Alhasanah, F., Sencaki, D. B., & Setyaningrum, N. (2025). Advancing Land Use Modeling with Rice Cropping Intensity: A Geospatial Study on the Shrinking Paddy Fields in Indonesia. *Geographies*, 5(3), 31. <https://doi.org/10.3390/geographies5030031>
- Gunawan, B., Abdoellah, O. S., Hadi, F., Alifi, G. J., Suhendi, R. N., Aisharya, I. Y., & Gunawan, W. (2023). From Laborers to Coffee Farmers: Collaborative Forest Management in West Java, Indonesia. *Sustainability*, 15(9), 7722. <https://doi.org/10.3390/su15097722>
- Hariyanto, A. D., Yudono, A., & Wicaksono, A. D. (2022). Land Cover Change Simulation Based on Cellular Automata Using Artificial Neural Network Model Transition in Kedungkandang District, Malang City. *International Conference on Indonesian Architecture and Planning*, 489–507. [https://doi.org/10.1007/978-981-99-1403-6\\_33](https://doi.org/10.1007/978-981-99-1403-6_33)
- Hariyanto, A. D., Yudono, A., & Wicaksono, A. D. (2024). Comparison of Land Cover Change Prediction Models: A Case Study in Kedungkandang District, Malang City. *Geoplanning: Journal of Geomatics and Planning*, 11(1), 85–98. <https://doi.org/10.14710/geoplanning.11.1.85-98>
- Lightfoot, D. R., & Butler, D. R. (1987). Recognition and assessment of error in geographic information systems. *Photogrammetric Engineering and Remote Sensing*, 53(10), 1423–1430. [https://www.academia.edu/download/90543545/1987\\_oct\\_1423-1430.pdf](https://www.academia.edu/download/90543545/1987_oct_1423-1430.pdf)
- MacEachren, A. M. (1992). Visualizing Uncertain Information. *Cartographic Perspectives*, 13, 10–19. <https://doi.org/10.14714/CP13.1000>
- Maling, D. H. (2013). *Measurements from maps: Principles and methods of cartometry*. Elsevier. [https://doi.org/10.1016/0160-9327\(89\)90096-3](https://doi.org/10.1016/0160-9327(89)90096-3)
- Manandhar, R., Odeh, I. O., & Ancev, T. (2009). Improving the Accuracy of Land Use and Land Cover Classification of Landsat Data Using Post-Classification Enhancement. *Remote Sensing*, 1(3), 330–344. <https://doi.org/10.3390/rs1030330>
- Nunes Machado, A., Costa, K., & Wachholz, F. (2024). **SENSORIAMENTO REMOTO APLICADO À PRÁTICA DE CAMPO NO ENSINO DE HIDROGRAFIA E GEOMORFOLOGIA FLUVIAL: UMA ANÁLISE DO ARQUIPÉLAGO DE ANAVILHANAS (AMAZONAS-BRASIL)**. In *Remote sensing and field practice in geography and geomorphology* (pp. 103–112). Editora Científica. ISBN: 978-85-5467-374-1
- Queiroga, F. M. (2020). *Interferência das Autoestradas na Conectividade Funcional da Paisagem para os Anfíbios do Peloponeso na Grécia* [Master's thesis, Universidade do Porto (Portugal)]. ISBN: 9798383454237
- Rayner, M. J. (2022). *Synergies and trade-offs in ecosystem services across the Upper Welland catchment* [PhD thesis, University of Leicester]. <https://doi.org/10.25392/leicester.data.21324360.v1>

- Rayner, M., Balzter, H., Jones, L., Whelan, M., & Stoate, C. (2021). Effects of improved land-cover mapping on predicted ecosystem service outcomes in a lowland river catchment. *Ecological Indicators*, 133, 108463. <https://doi.org/10.1016/j.ecolind.2021.108463>
- Rayner, M., Whelan, M., Stoate, C., Szczur, J., Balzter, H., & Jones, L. (2021). Historical Declines in Semi-Natural Habitat Across an Agricultural Landscape Drive Modelled Losses in Wild Bee Abundance. *SSRN*. <https://doi.org/10.2139/ssrn.3969794>
- Schowengerdt, R. A. (2007). CHAPTER 9 - Thematic Classification. In R. A. Schowengerdt (Ed.), *Remote Sensing (Third Edition)* (Third Edition, pp. 387–XXXIII). Academic Press. <https://doi.org/10.1016/B978-012369407-2/50012-7>
- Senterre, B., Mederic, E., & Berke, G. (2023). *Flora and vegetation of Assomption Island (Seychelles)* (Vol. 775). Island Conservation Society (ICS), Pointe Larue, PO Box. [https://www.researchgate.net/profile/Bruno-Senterre/publication/378943939\\_Flora\\_and\\_vegetation\\_of\\_Assomption\\_Island\\_Seychelles/links/65f2f44fc05fd2688010af87/Flora-and-vegetation-of-Assomption-Island-Seychelles.pdf](https://www.researchgate.net/profile/Bruno-Senterre/publication/378943939_Flora_and_vegetation_of_Assomption_Island_Seychelles/links/65f2f44fc05fd2688010af87/Flora-and-vegetation-of-Assomption-Island-Seychelles.pdf)
- Thakkar, A. K., Desai, V. R., Patel, A., & Potdar, M. B. (2017). Post-classification corrections in improving the classification of Land Use/Land Cover of arid region using RS and GIS: The case of Arjuni watershed, Gujarat, India. *The Egyptian Journal of Remote Sensing and Space Science*, 20(1), 79–89. <https://doi.org/10.1016/j.ejrs.2016.11.006>
- Vallet, A., Dupuy, S., Verlynde, M., & Gaetano, R. (2024). Generating high-resolution land use and land cover maps for the greater Mariño watershed in 2019 with machine learning. *Scientific Data*, 11(1), 915. <https://doi.org/10.1038/s41597-024-03750-x>