








scarlet2: Astronomical scene modeling in JAX

Peter Melchior ¹, Charlotte Ward ^{1,2}, Benjamin Remy ^{1,3}, Matt L. Wiemann ¹, and Jared Siegel ¹

¹ Princeton University, United States  ² Pennsylvania State University, United States  ³ University of Chicago, United States   Corresponding author

DOI: [10.21105/joss.09646](https://doi.org/10.21105/joss.09646)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Ivelina Momcheva](#)  

Reviewers:

- [@AlexandreAdam](#)
- [@Jammy2211](#)

Submitted: 29 July 2025

Published: 29 April 2026

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Summary

Large astronomical imaging surveys now contain billions of celestial sources, from stars in our Galaxy to galaxies towards the edge of the observable Universe. Because of improvements in resolution and sensitivity of current and future observing instruments, the images reveal more information than ever before. But they are also more difficult to analyze because galaxies exhibit complex morphologies, which cannot be described by traditional parametric models. And because there are so many sources, they routinely overlap with each other, either due to physical interactions or due to their close alignment along the line of sight. To extract all information of interest and avoid biases from incorrect modeling assumptions, it is therefore necessary to simultaneously model full scenes comprising many sources instead of analyzing each source separately, and each of the source models may itself need to be composed of multiple, morphologically complex components.

Statement of need

scarlet2 is a Python package for full-scene modeling in observational astronomy. It inherits modeling assumptions from scarlet ([Melchior et al., 2018](#)), namely that a scene comprises multiple sources, each source comprises multiple components, and each component is determined by a spectrum model and a morphology model, whose outer product represents the light emission in a sky region as a hyperspectral data cube (wavelength \times height \times width). scarlet2 retains the object-oriented paradigm and many classes and functions from scarlet, but augments standard Python with the JAX library ([Bradbury et al., 2018](#)) and the Equinox package ([Kidger & Garcia, 2021](#)) for automatic differentiation and just-in-time compilation.

scarlet2 acts as a flexible, modular, and extendable modeling language for celestial sources that combines parametric and non-parametric models to describe complex scenarios such as multi-source blending, strong-lensing systems, supernovae and their host galaxies, etc. As a modeling language, scarlet2 is agnostic about the optimization or inference method the user wants to employ, but it also provides methods to optimize the likelihood function or sample from the posterior, which utilize the Optax package ([DeepMind et al., 2020](#)) or the NumPyro inference framework ([Bingham et al., 2019](#); [Phan et al., 2019](#)), respectively. The likelihood of multiple observations (at different resolutions, wavelengths, or observing epochs) can be combined for a joint model of static and transient sources. To match the coordinates from different observations, scarlet2 utilizes the Astropy package ([Astropy Collaboration, 2013](#)). scarlet2 can also interface with deep learning methods. Besides being natively portable to GPUs, parameters can be specified with neural networks as data-driven priors, which helps break the degeneracies that arise when multiple components are fit simultaneously ([Sampson et al., 2024](#)).




Figure 1: Scene with seven detected sources in multi-band images from the Hyper Suprime-Cam Subaru Strategic Program. Each source is modelled with a non-parametric spectrum and morphology (first panel), the entire scene is then convolved with the telescope's point spread function (second panel) and compared to the observations (third panel). The residuals (fourth panel) reveal the presence of previously undetected sources and source components (e.g. in the center of source #1).

To support the wide range of scientific studies that will be made with large sky surveys, `scarlet2` was designed with flexibility and ease of use in mind. Several publications have developed and demonstrated its capabilities, including modeling of interstellar dust embedded in distant galaxies (Siegel & Melchior, 2025) and of transient sources such as active galactic nuclei (Ward et al., 2025) and tidal disruption events (Yao et al., 2025). We have recently added support for users to make effective modeling choices and to validate their inference results. Future developments will integrate `scarlet2` into cloud-based science platforms and create a robust processing pipeline for joint pixel-level analyses of surveys from the Vera C. Rubin Observatory, the Euclid mission, the Nancy Grace Roman Space Telescope, and the La Silla Schmidt Southern Survey.

State of the field

`GALFIT` (Peng et al., 2002) implements parametric Sérsic model fitting in single-band, single-epoch data. The underlying algorithms are closed source and released only as precompiled binaries. `GALAPAGOS` (Barden et al., 2012) provides wrappers around `GALFIT` in the proprietary IDL programming language; `GALAPAGOS-2` (Häußler et al., 2013) added multi-band fitting capabilities. The `Tractor` (Lang et al., 2016) is implemented in Python and allows modeling galaxies in multi-band imaging at different resolutions as well as posterior sampling, but remains limited to parametric source profiles. `AstroPhot` (Stone et al., 2023) offers an astronomical modeling framework based on PyTorch (Paszke et al., 2019), which allows one to optimize multiple sources with different parametric models to multiple images at different resolutions, and includes posterior sampling techniques. None of the implementations above support fully non-parametric galaxy models or neural network priors.

Acknowledgements

We acknowledge contributions from the [LINCC Frameworks Incubator Program](#), in particular from software engineers Max West, Drew Oldag, and Sean McGuire, in adopting comprehensive software workflows through the Python Project Template (Oldag et al., 2024) and creating a user-focused recommendation and validation suite.

In addition to dependencies mentioned earlier, `scarlet2` makes use of NumPy (Harris et al., 2020), Matplotlib (Hunter, 2007), and h5py (Collette, 2013).

References

- Astropy Collaboration. (2013). Astropy: A community Python package for astronomy. *Astronomy and Astrophysics*, 558, A33. <https://doi.org/10.1051/0004-6361/201322068>
- Barden, M., Häußler, B., Peng, C. Y., McIntosh, D. H., & Guo, Y. (2012). GALAPAGOS: from pixels to parameters. *Monthly Notices of the Royal Astronomical Society*, 422(1), 449–468. <https://doi.org/10.1111/j.1365-2966.2012.20619.x>
- Bingham, E., Chen, J. P., Jankowiak, M., Obermeyer, F., Pradhan, N., Karaletsos, T., Singh, R., Szerlip, P. A., Horsfall, P., & Goodman, N. D. (2019). Pyro: Deep universal probabilistic programming. *Journal of Machine Learning Research*, 20, 28:1–28:6. <http://jmlr.org/papers/v20/18-403.html>
- Bradbury, J., Frostig, R., Hawkins, P., Johnson, M. J., Leary, C., Maclaurin, D., Necula, G., Paszke, A., VanderPlas, J., Wanderman-Milne, S., & Zhang, Q. (2018). JAX: Composable transformations of Python+NumPy programs (Version 0.3.13). <http://github.com/google/jax>
- Collette, A. (2013). *Python and HDF5*. O'Reilly.
- DeepMind, Babuschkin, I., Baumli, K., Bell, A., Bhupatiraju, S., Bruce, J., Buchlovsky, P., Budden, D., Cai, T., Clark, A., Danihelka, I., Dedieu, A., Fantacci, C., Godwin, J., Jones, C., Hemsley, R., Hennigan, T., Hessel, M., Hou, S., ... Viola, F. (2020). *The DeepMind JAX Ecosystem*. <http://github.com/google-deepmind>
- Harris, C. R., Millman, K. J., Walt, S. J. van der, Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., Kerkwijk, M. H. van, Brett, M., Haldane, A., Río, J. F. del, Wiebe, M., Peterson, P., ... Oliphant, T. E. (2020). Array programming with NumPy. *Nature*, 585(7825), 357–362. <https://doi.org/10.1038/s41586-020-2649-2>
- Häußler, B., Bamford, S. P., Vika, M., Rojas, A. L., Barden, M., Kelvin, L. S., Alpaslan, M., Robotham, A. S. G., Driver, S. P., Baldry, I. K., Brough, S., Hopkins, A. M., Liske, J., Nichol, R. C., Popescu, C. C., & Tuffs, R. J. (2013). MegaMorph - multiwavelength measurement of galaxy structure: complete Sérsic profile information from modern surveys. *Monthly Notices of the Royal Astronomical Society*, 430(1), 330–369. <https://doi.org/10.1093/mnras/sts633>
- Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. *Computing in Science & Engineering*, 9(3), 90–95. <https://doi.org/10.1109/MCSE.2007.55>
- Kidger, P., & Garcia, C. (2021). Equinox: Neural networks in JAX via callable PyTrees and filtered transformations. *Differentiable Programming Workshop at Neural Information Processing Systems 2021*. <https://doi.org/10.48550/arXiv.2111.00254>
- Lang, D., Hogg, D. W., & Mykytyn, D. (2016). *The Tractor: Probabilistic astronomical source detection and measurement*. Astrophysics Source Code Library, record ascl:1604.008.
- Melchior, P., Moolekamp, F., Jerdee, M., Armstrong, R., Sun, A.-L., Bosch, J., & Lupton, R. (2018). SCARLET: Source separation in multi-band images by Constrained Matrix Factorization. *Astronomy and Computing*, 24, 129. <https://doi.org/10.1016/j.ascom.2018.07.001>
- Oldag, D., DeLucchi, M., Beebe, W., Branton, D., Campos, S., Chandler, C. O., Christofferson, C., Connolly, A., Kubica, J., Lynn, O., Malanchev, K., Malz, A. I., Mandelbaum, R., McGuire, S., & Wenneman, C. (2024). A Python project template for healthy scientific software. *Research Notes of the American Astronomical Society*, 8(5), 141. <https://doi.org/10.3847/2515-5172/ad4da1>
- Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., Killeen, T., Lin, Z.,

- Gimelshein, N., Antiga, L., Desmaison, A., Kopf, A., Yang, E., DeVito, Z., Raison, M., Tejani, A., Chilamkurthy, S., Steiner, B., Fang, L., ... Chintala, S. (2019). PyTorch: An imperative style, high-performance deep learning library. In *Advances in neural information processing systems 32* (pp. 8024–8035). Curran Associates, Inc. <http://papers.neurips.cc/paper/9015-pytorch-an-imperative-style-high-performance-deep-learning-library.pdf>
- Peng, C. Y., Ho, L. C., Impey, C. D., & Rix, H.-W. (2002). Detailed structural decomposition of galaxy images. *The Astronomical Journal*, *124*(1), 266–293. <https://doi.org/10.1086/340952>
- Phan, D., Pradhan, N., & Jankowiak, M. (2019). Composable effects for flexible and accelerated probabilistic programming in NumPyro. *arXiv Preprint arXiv:1912.11554*. <https://doi.org/10.48550/arXiv.1912.11554>
- Sampson, M. L., Melchior, P., Ward, C., & Birmingham, S. (2024). Score-matching neural networks for improved multi-band source separation. *Astronomy and Computing*, *49*, 100875. <https://doi.org/10.1016/j.ascom.2024.100875>
- Siegel, J. C., & Melchior, P. (2025). Spatially resolved galaxy–dust modeling with coupled data-driven priors. *The Astrophysical Journal*, *986*(2), 212. <https://doi.org/10.3847/1538-4357/add3f9>
- Stone, C. J., Courteau, S., Cuillandre, J.-C., Hezaveh, Y., Perreault-Levasseur, L., & Arora, N. (2023). AstroPhot: Fitting everything everywhere all at once in astronomical images. *Monthly Notices of the Royal Astronomical Society*, *525*(4), 6377–6393. <https://doi.org/10.1093/mnras/stad2477>
- Ward, C., Melchior, P., Sampson, M. L., Burke, C. J., Siegel, J., Remy, B., Birmingham, S., Ramey, E., & Velzen, S. van. (2025). Disentangling transients and their host galaxies with scarlet2: A framework to forward model multi-epoch imaging. *Astronomy and Computing*, *51*, 100930. <https://doi.org/10.1016/j.ascom.2025.100930>
- Yao, Y., Chornock, R., Ward, C., Hammerstein, E., Sfaradi, I., Margutti, R., Kelley, L. Z., Lu, W., Liu, C., Wise, J., Sollerman, J., Alexander, K. D., Bellm, E. C., Drake, A. J., Fremling, C., Gilfanov, M., Graham, M. J., Groom, S. L., Hinds, K. R., ... Wold, A. (2025). A massive black hole 0.8 kpc from the host nucleus revealed by the offset tidal disruption event AT2024tvd. *The Astrophysical Journal Letters*, *985*(2), L48. <https://doi.org/10.3847/2041-8213/add7de>